

A Volunteer's Handbook for Monitoring New Hampshire's Salt Marshes



A Volunteer's Handbook for Monitoring New Hampshire Salt Marshes



Prepared By:

Jen Drociak
New Hampshire Coastal Program

&

Grace Bottitta
Ducks Unlimited

Through funding granted in part from the Gulf of Maine Council on the Marine Environment and the National Oceanic & Atmospheric Administration (NOAA) Community-Based Restoration Program awards # NA17021529 and NA17FZ1390.



Printed on Recycled Paper
May 2003
Revised: May 2004, January 2005, September 2005

Sponsoring Organizations

NH Department of Environmental Services

NH Coastal Program

50 International Drive – Suite 200

Portsmouth, NH 03801

(603) 559-1500

www.des.nh.gov/coastal



Ducks Unlimited, Inc.

Atlantic Coast Ecosystem Initiative

331 Metty Drive, Suite 4

Ann Arbor, MI 48103

(734) 623-2000

(603) 778-0704

www.ducks.org/conservation



University of New Hampshire

Jackson Estuarine Laboratory

85 Adams Point Road

Durham, NH 03824

(603) 862-2175

<http://marine.unh.edu/jel/jel.html>



Association of the US Delegates to the Gulf of Maine Council

PO Box 2246

South Portland, ME

(207) 799-7474

www.gulfofmaine.org



National Oceanic and Atmospheric Administration (NOAA)

Community-Based Restoration Programs

Restoration Center

1315 East-West Highway

Silver Spring, Maryland 20910

(301) 713-0174

www.nmfs.noaa.gov/habitat/restoration

Acknowledgements

This manual is an integration of protocols found in the “Regional Standards to Identify and Evaluate Tidal Wetland Restoration in the Gulf of Maine” (*Global Programme of Action Coalition for the Gulf of Maine Workshop, 1999*), “A Volunteer’s Handbook for Monitoring New England Salt Marshes” (*Massachusetts Office of Coastal Zone Management and the Executive Office of Environmental Affairs, 2002*), and “Field Methods Manual: U.S. Fish and Wildlife Service (Region 5) Salt Marsh Study” (*U.S. Fish and Wildlife Service, 2002*).

- ❖ **Susan Adamowicz:** U.S. Fish and Wildlife Service
- ❖ **Bruce Carlisle:** Massachusetts Office of Coastal Zone Management
- ❖ **Michelle Dionne:** Wells National Estuarine Research Reserve
- ❖ **Anne M. Donovan:** Massachusetts Office of Coastal Zone Management
- ❖ **Anna L. Hicks:** Independent Consultant
- ❖ **Vivian S. Kooken:** Salem Sound 2000
- ❖ **Hilary Neckles:** U.S. Geological Survey
- ❖ **Kenneth B. Raposa:** University of Rhode Island
- ❖ **Charles T. Roman:** U.S. Geological Survey
- ❖ **Gregory Shriver:** National Park Service
- ❖ **Jan P. Smith:** Massachusetts Bays National Estuary Program
- ❖ **Jan Taylor:** U.S. Fish and Wildlife Service
- ❖ **Anthony R. Wilbur:** Massachusetts Office of Coastal Zone

Content Contributions

- ❖ **Dave Burdick:** University of New Hampshire, Jackson Estuarine Laboratory (2003, 2004)
- ❖ **Matt Craig:** Ducks Unlimited, Inc. (2004)
- ❖ **Stephanie Wojtowicz:** Ducks Unlimited, Inc. and Antioch New England Graduate School (2004)
- ❖ **Lois Winter:** U.S. Fish and Wildlife Service, Gulf of Maine Program (2004)

Additional Contributions

- ❖ **Brandi Bornt:** Ducks Unlimited, Inc. and Antioch New England Graduate School (2005)
- ❖ **Ted Diers:** New Hampshire Coastal Program (2003)
- ❖ **Alyson Eberhardt:** University of NH, Jackson Estuarine Laboratory (2004)
- ❖ **Pat Gruttemeyer:** New Hampshire Department of Environmental Services (2004)
- ❖ **Sean McDermott:** National Oceanic and Atmospheric Administration (2003)
- ❖ **Ann Reid:** Great Bay Coast Watch (2003)




Table of Contents

	Page
Sponsoring Organizations	i
Acknowledgements	ii
New Hampshire Coastal Zone Map	iii
1. An Introduction to Salt Marsh Monitoring.....	1
2. New England Salt Marshes: Ecology, Importance, and Conservation.....	7
3. Elements of Study Design.....	15
4. Data Management, Quality Assurance, and Quality Control.....	21
5. Ecology of Salt Marsh Vegetation.....	23
6. Nekton.....	31
7. Birds.....	37
8. Soil Characteristics	45
9. Hydrology.....	49

Appendices

A. Suggested Reading.....	53
B. Glossary.....	57
C. Lift Net & Ditch Net Materials and Construction.....	61
D. Field Data Sheets.....	65

A Note About This Publication

This manual was created to provide volunteers and project collaborators a brief overview of the ecology and importance of local salt marshes and the human activities that have impacted them. Specific protocols will be used to monitor a few indicators of salt marsh functions to facilitate management decisions. For further information and resources on these topics please refer to Appendix A.

Terms printed in bold type at their first appearance are defined in the Glossary (Appendix B).

If you believe a particular salt marsh or other coastal wetland ecosystem is in need of restoration, please contact the New Hampshire Coastal Program at (603) 559-1500.

Chapter 1: An Introduction to Salt Marsh Monitoring

Salt marshes are beautiful coastal landscapes that provide rich **habitat** to a great **diversity** of plants, **invertebrates**, fish, birds, and mammals. For many people, the opportunity to see snowy egrets stalking fish is reason enough to be concerned about salt marsh health. Salt marshes are extremely important for a variety of reasons besides their beauty or the biological diversity they support. Salt marshes are among the most productive ecosystems on the planet, and serve as nursery grounds for several economically important fish and shellfish species. Salt marshes also help fuel **food webs** by recycling and exporting tremendous amounts of nutrients. Salt marshes protect shorelines from storm damage by dispersing wave and tide energy, and help purify water by assimilating potential pollutants.



*A great blue heron forages for food.
Photo by UNH*

Over the last three centuries, vast areas of salt marshes have been ditched, drained, and filled by humans who perceived them as barren unproductive areas of little economic value.

Direct **wetland** filling, point source pollution, nonpoint source pollution, invasive species, ditching, and restriction of tide flow by road and railroad crossings have all taken a heavy toll on New England coastal salt marshes. In 1969, John and Mildred Teal published their book *Life and Death of a Salt Marsh*, which highlighted the beauty, importance, and plight of these ecosystems. This book helped foster public appreciation for salt marshes and launched the next three decades of salt marsh conservation.



*Pickering Brook saltmarsh, Greenland, NH.
Photo by GBCW*

Scientists use a variety of methods to assess salt marsh health, also referred to as **biological integrity**. By comparing modern aerial photographs and maps with historical documents using geographic information systems (GIS), researchers are able to estimate changes to wetland area over time. Generally, these studies have found a decline in salt marsh area resulting from human manipulation and development. Researchers also collect field measurements of various biological, physical, and chemical **parameters**, which are used as **indicators** of overall salt marsh health.

Although regulations help reduce further human impacts to salt marshes, they are not sufficient to adequately protect these critical and sensitive habitats. Scientists are currently developing and employing tools to detect **biological impairment** in salt marsh ecosystems. The information generated from these assessments will help to improve regulatory and other protection efforts.

A growing number of organizations are becoming involved in conservation projects aimed at restoring or protecting salt marshes. Through the New Hampshire Marsh Monitoring Program (NHMMP), local

volunteers, under the guidance of salt marsh experts, are collecting field data to document salt marsh health and look for evidence of habitat degradation and biological impairment. Scientists refer to this effort as **monitoring**, or “the unbiased collection and precise recording of data over time.” The NHMMP applies an integrated approach to monitoring that combines biological, physical, and chemical measurements. Use of this approach provides a comprehensive overview of salt marsh ecosystem health. By assisting with monitoring efforts, volunteers provide a valuable service to scientists trying to develop a better understanding of salt marshes, to managers aiming to protect and restore salt marshes, and to their communities.

Salt Marsh Monitoring in New England

The protocols contained in this manual are the culmination of over ten years of collaborative effort among wetland scientists throughout the Northeast, including representatives from state and federal agencies and non-profit groups. The goal was to develop salt marsh bioassessment techniques for measuring wetland health by examining resident plants, animals, and their habitat. Despite decades of research into salt marsh ecological processes, widespread bioassessment of New England salt marshes didn't occur until 1995 following a series of pilot projects to develop monitoring protocols. In 1997, the U.S. Environmental Protection Agency declared wetland monitoring a national priority and convened the national Biological Assessment of Wetlands Workgroup in which wetland scientists from federal and state agencies and universities collaborated to improve methods for evaluating the biological integrity of wetlands. In 1999, the Global Programme of Action Coalition (GPAC) held a workshop for resource managers and scientists to recommend regional standards for salt marsh monitoring protocols. Several restoration partnerships in Maine have used contractors instead of volunteers to collect pre and post restoration information. The methods used by the New Hampshire Marsh Monitoring Program and described in this manual are volunteer-friendly and are consistent with recommendations from the GPAC report, entitled *Regional Standards to Identify and Evaluate Tidal Restoration in the Gulf of Maine* and the *Field Methods Manual: U.S. Fish and Wildlife Service (Region 5) Salt Marsh Study* used in several federal wildlife refuges and coastal national parks along the North Atlantic coast.

The Importance of Volunteer Monitoring

Volunteer participation in salt marsh monitoring efforts provides an invaluable service to resource managers, scientists, and communities throughout New England. Volunteer participation enables scientists to develop larger data sets, provides resource managers with access to increased federal funding for protection and restoration efforts, and provides program coordinators with valuable feedback about data collection and recording methods. Coastal resource managers rely on current data to assess and identify

potential threats to salt marsh health in order to develop effective protection and restoration strategies. Scientists who assess the effectiveness of various restoration and protection strategies incorporate data into a regional database. Thus, by participating in the New Hampshire Marsh Monitoring Program, volunteers provide direct support not only to protection and restoration efforts in the state but also to similar efforts throughout the Gulf of Maine.



Volunteer monitors also become informed ambassadors for salt marshes by being active in local planning and decision-making and fostering salt marsh stewardship and awareness in coastal communities. Volunteers, communities, resource managers, and scientists all benefit from this type of partnership.

The Role of Volunteer Monitors

Some people may view participation in a volunteer monitoring program as a daunting task and wonder “What can I do?” “How can I help?” and “What will my contribution mean?” As participants in the New Hampshire Marsh Monitors Program, volunteers will receive training to collect data in salt marshes throughout the seacoast area. Once training has been completed, volunteers will join scientists, resource managers, and other experts in the marshes to collect and record field data. You do not need a college degree in biology to be a volunteer monitor – all you need is enthusiasm and a willingness to learn. Participants will have the opportunity to adopt regular monitoring schedules at particular sites, if interested, or may sign up for pre-determined monitoring sessions spread intermittently throughout the field season. Program coordinators will assist volunteers with scheduling, data collection, and data recording, and will track volunteer participation as match for federal funds.

The Application of Volunteer Data

For decades volunteers have been involved in monitoring programs by counting birds, taking water quality samples in lakes, listening for breeding amphibians, and collecting stream invertebrates to provide valuable data to state and federal agencies.

Many scientists and resource managers have invested time and resources to train volunteers to conduct salt marsh monitoring. This is necessary so that volunteers can collect data that are as accurate and defensible as data collected by staff scientists. The guidelines and procedures outlined in this manual and taught at workshops are not mere suggestions – volunteers need to follow these guidelines explicitly to ensure data quality. Using this manual, volunteers can gather data that will directly influence the conservation and management of coastal resources.



Nekton sampling using a lift net. Photo by DU

Goals of this Manual

- ❖ Provide volunteer monitors with a brief background and training necessary to successfully collect scientifically reliable field data.
- ❖ Foster salt marsh education and stewardship in communities throughout the seacoast area.
- ❖ Promote and assist with the stewardship, restoration, and protection of salt marshes.
- ❖ Expand the number of qualified individuals who can help scientists study the condition of the region’s marshes.

Level of Technical Expertise

Volunteer monitoring requires only an interest in salt marshes, an ability to carefully navigate across sometimes-unpredictable marsh terrain, and a willingness to devote time and energy toward marsh conservation. This manual employs technical language and scientific species' names in order to communicate in well defined and commonly agreed upon scientific terminology. Volunteers are not expected to know and use all of these terms. However, volunteers are encouraged to learn scientific terminology and ask questions of program coordinators at any time throughout the field season.

Care of the Salt Marsh

Salt marshes are fragile and sensitive ecosystems. Volunteers should be mindful of how their activities affect the marsh and take appropriate steps to minimize impacts, since most types of monitoring require that volunteers walk across the marsh surface. Vegetation is easily trampled, and substrate erosion can be problematic. If possible, volunteers should minimize unnecessary trampling and follow paths at sites that they visit repeatedly. When entering estuarine streams and crossing ditches, select areas of bank that aren't too high or too steep; clambering up and down steep stream banks will quickly result in bank erosion.

Wildlife disturbance can be a concern at some locations. Birds may breed or nest in the salt marsh, marsh border, or adjacent dune areas. If possible, identify important breeding territories and avoid these areas during the nesting season. **Invasive species** such as the **non-indigenous** form of *Phragmites australis*, have become a major problem in coastal wetlands. Most invasive plants have evolved rapid dispersal abilities and rely on people and other animals to transport them to new sites. We can help curtail the spread of invasive species and pathogens by thoroughly rinsing waders, footwear, and sampling equipment immediately after leaving one marsh and before moving to another.

Safety Issues

Humidity in salt marshes can reach uncomfortable levels because of evaporation from saturated soils and transpiration from vegetation. The marsh can get very hot because there is no shade. In addition, light intensity is high and the marsh and surface water reflects sunlight. This lack of shade, high humidity, and continuous activity can cause monitors to get very hot. All volunteers should make sure to *bring plenty of water and sunscreen*.


Volunteers will be walking and wading in salt marshes. This can be an enjoyable experience, but volunteers should take steps to protect themselves and the salt marsh from harm. Salt marshes can be very difficult to walk through. Volunteers must be prepared for all types of conditions. Scorching sun, biting flies, ticks, poison ivy, hip-deep mud, slippery surfaces, and unstable footing can combine to make an uncomfortable experience for unprepared volunteers. *Step carefully!*

Estuarine streams and tidal flats contain deceptively deep mud, and when you are stuck knee deep as the tide rolls in there is nothing more welcomed than a helping hand from a fellow team member. Marshes also have a large appetite for loose-fitting shoes! It is easier to sink your foot into deep mud than it is to pull your foot out, and shoes are often lost if they are not laced tightly. The best types of footwear are *knee-high rubber boots* or hip waders.

Poison ivy is common in the high marsh-upland transitional zone, and it is important that sensitive individuals wear long clothing to protect themselves. In addition, ticks, mosquitoes, and other biting insects can be both a nuisance and serious health threat, carrying West Nile virus, Lyme disease, and other maladies. Long clothing and insect repellent are effective deterrents, and volunteers should thoroughly check themselves for ticks after leaving a wetland.

Essential Field Equipment

Volunteers should always bring the following items when entering a marsh to ensure that they will be comfortable and safe:



Chapter 2: New England Salt Marshes Ecology, Value, and Conservation



Little River, North Hampton, NH. Photo by NHCP

Salt marshes are among the most productive natural ecosystems on Earth. They support a rich diversity of plants and animals that are uniquely adapted to inhabit wetlands that constantly change with the ebb and flow of the tide. Today, salt marshes are treasured for the biodiversity they support, their contribution to marine productivity and commercial fisheries, their ability to anchor sediments and protect shorelines from erosion and flood damage, and their capacity to improve water quality.

Despite their importance, humans have impacted New England salt marshes since the Colonial period. Marshes were ditched to promote the growth of salt hay, *Spartina patens*, which was then fed to livestock. In the early to mid 20th Century, salt marshes were perceived as barren, unproductive, mosquito-filled wastelands that could only be improved by filling, draining, or dredging. As a result, countless acres of salt marshes have been converted to roads, farms, parking lots, neighborhoods, and even cities. For instance, large areas of Boston were once productive salt marshes:

“There are at present about 3,240 acres of city real estate in an area that contains old Boston, Roxbury and Back Bay...When the Puritans arrived to settle this area, there existed only 1,185 acres of dry land on which to build. Four hundred eighty-five acres of the present 3,240 acres were salt marsh and 1,570 acres were shallow water, which was part marsh, part mud and sand flat, and part open water even at low tide. There was a gain of 2,055 acres of dry land made by filling the marshes and lowlands.” John and Mildred Teal 1969, *Life and Death of the Salt Marsh*.

Our understanding of the importance of salt marshes has increased in the last few decades, and today there are strict laws and regulations designed to protect salt marshes from despoliation. Yet, coastal development continues throughout New England and natural resources face increasing pressure as communities try to find a balance between development and healthy natural ecosystems.

This chapter introduces some important concepts of salt marsh biology and ecology, including their formation and



Vol's Island, Newmarket, NH.

Photo by Great Bay Resource Protection Partnership

succession, characteristic plant communities, food webs, and importance as nursery area and wildlife habitat. This chapter also covers why salt marshes are important to humans, and the many ways that humans have destroyed, degraded, and polluted these valuable ecosystems. An understanding of salt marsh biology and ecology is imperative for anyone who is conducting environmental impact studies or basic monitoring in these habitats. The current condition of salt marshes is the product of ever changing natural processes and **human disturbances**, and any successful monitoring program needs to be mindful of both.

Biology and Ecology of New England Salt Marshes

Life in coastal wetlands is characterized by extreme fluctuations in salinity, saturation, and weather on a daily and seasonal basis. In general, New England coastal wetlands experience regular patterns of flooding and exposure with the ebb and flow of the tide. In a single day, a crab in a coastal wetland might be submerged in seawater, caught in a rainstorm, exposed to the open atmosphere, and as a result, experience salinities between 10 to 35 parts per thousand (ppt). Normal seawater in the Gulf of Maine ranges from salinities of 30 to 35 ppt. Plants and animals inhabiting these habitats must be highly specialized to deal with such extremes.

Salt Marsh Development

Coastal wetlands exist in areas that are periodically flooded by tidal waters. They develop along embayments, barrier beaches, islands, and **estuaries** that form the link between the ocean and non-tidal freshwater habitats. While this manual focuses on salt marshes, tidal mud flats and rocky beaches are examples of the other coastal wetlands whose conditions are less hospitable to the establishment of plant communities.

Salt marshes develop in sheltered coastal areas where the absence of severe winds and waves allows fine sediments to settle and accumulate and for plants to eventually take root. Salt marshes form along barrier landforms and islands, coastal ponds, and tidal creeks or rivers. Salt marshes are common along the Atlantic seaboard as far north as mid-coast Maine.



Two other types of marshes are influenced by tides yet support different plant communities. Brackish marshes exist further inland along estuarine systems and have salinities ranging from 0.5 to 18 ppt. Brackish marshes can support typical salt marsh plants along the seaward edge of the marsh, but also support a high diversity of freshwater and slightly salt-tolerant wetland plants. Tidal freshwater marshes occur at the inland limit of estuaries where tides continue to cause fluctuating water levels but seawater fails to penetrate. Tidal freshwater marshes support a high diversity of wetland plants that are intolerant of salinity. Being able to identify plant and animal species from brackish and freshwater marshes is an important component to studying salt marshes because tide restrictions frequently cut off salt marshes from their tidal influence, causing salt **tolerant** organisms to be out-competed and replaced by brackish and freshwater organisms.

Salt Marsh Succession

Salt marshes develop in sheltered coastal areas that are protected from severe wind and wave action. The combination of low energy and deposition of fine sediments (sand and silt) favors the establishment of plant communities. One of the first plants to take hold in these areas is *Spartina alterniflora*, whose seeds are dispersed by wind and water. *Spartina alterniflora* is a perennial plant that develops an extensive root system, called **rhizomes**, which stabilize sediments and reduce erosion. As this plant establishes itself, it forms dense stands that buffer wave energy and trap sediments, promoting further development of the infant salt marsh.

Bacteria and fungi slowly decay organic matter trapped by a growing stand of *Spartina alterniflora*. Over time, the accumulation of dead and decaying matter results in the formation of peat. Peat accumulation occurs for many years and raises the elevation of the marsh enough to reduce flooding frequency. Once this occurs, high marsh plants such as *Spartina patens* and *Distichlis spicata* can become established, which in turn accelerates peat accumulation and eventually allows a greater diversity of salt marsh plants to exist there. A mature salt marsh has a well-defined low marsh and high marsh that continue to expand seaward and landward over time. The landward migration of the salt marsh occurs as the marsh keeps pace with sea level rise. As the height of the sea increases very gradually, so too does the surface of the marsh. Unfortunately, coastal development at the marsh's edge prohibits this landward migration, and over a long time, sea level rise and hardened shorelines may become a major cause of salt marsh loss.

Higher than normal tides deposit large amounts of dead plant material or other debris on the marsh, creating bare or open areas by shading and killing the plants below and often slightly lowering the marsh elevation underneath. These depressions may become pannes or pools. Pioneer salt-tolerant plants like *Salicornia spp.* and *Sueda linearis* quickly colonize these open areas. Over time, typical high marsh plants may outcompete and replace opportunistic species. The important thing to remember about salt marsh succession is that the appearance, productivity, and biological diversity of a salt marsh constantly change due to natural processes. Human disturbance and pollution certainly affect salt marshes, but the distinction between natural processes and human impacts is sometimes fuzzy, and can lead to uncertainty in environmental impact studies.



Sueda linearis, a pioneer plant, colonizes a salt marsh.
Photo by DU

Salt marshes are detritus-based ecosystems. Scientists define **detritus** as decomposing plant and animal material. Bacteria, fungi, microscopic plants and animals, and larger animals all contribute to the decomposition of dead plants, resulting in detritus. Many invertebrates eat detritus and associated decomposers. These invertebrates are called detritivores. Certain detritivores eat detritus directly (snails, crabs, and amphipods), whereas others eat sediment that contains detritus (fiddler crabs, snails, shrimp, and worms).

A third group of consumers are filter feeders, which include clams, mussels, and some worms. Filter feeders actively filter food (phytoplankton, zooplankton, detritus) from the water column. A fourth group of consumers are predators, which includes a wide variety of invertebrate and vertebrate animals. Terrestrial insects and spiders are important predators in the high marsh vegetation. Mummichogs, grass shrimp, and blue crabs are three common predators in the low marsh. There are also many birds that prey upon salt marsh animals, such as terns, plovers, egrets, and herons.

Importance of New England Salt Marshes

Salt marshes are dynamic and productive ecosystems that provide important benefits to humans and wildlife alike. Most notably, salt marshes are important nursery grounds and wildlife habitat, provide flood and erosion control, improve water quality, and offer recreational retreats.

Nursery Grounds and Wildlife Habitat

A rich diversity of invertebrate and vertebrate animals depends on salt marshes during some stage of their life cycle. Many invertebrates are well adapted to daily cycles of exposure and flooding associated with the change of tides, providing these organisms with competitive advantages in a productive ecosystem.



Herring. Photo by NMFS

Many animals are not adapted to live in salt marshes all the time, yet have found ways to capitalize on the food and safety that salt marshes provide during essential times of their life cycle. Many marine fish use salt marshes as breeding grounds or nursery habitats as juveniles, where they find an abundant supply of prey (such as worms, mollusks, and crustaceans). Menhaden, flounder, sea trout, spot, and striped bass are just a few examples of game fish that use salt marshes at some point during their lives. Non-game fish such as killifish and mummichogs also rely on salt marshes and are key forage species for game fish such as striped bass and bluefish. Blue crabs forage for prey in the low marsh during high tide but move offshore during low tide; unlike green crabs and fiddler crabs, blue crabs cannot tolerate long periods of exposure.

Animals do not need to enter salt marshes to benefit from their productivity. Every ebb tide carries a flush of nutrients and detritus into offshore areas, where it is eaten by a variety of consumers and helps fuel marine food webs. In addition, fish and crustaceans that feed in salt marshes and then move into offshore areas are essentially transferring salt marsh-derived nutrients into marine food webs.

Many birds use salt marshes to nest, breed, feed, or rest during migration. Some of these birds are rare and protected, such as the Northern Harrier, Least Tern, Osprey, and Saltmarsh Sparrow. Salt marshes are also critical habitat for the New Hampshire listed species such as the shortnose sturgeon. Deer, muskrats, otters, foxes, and coyotes may also forage in or near salt marshes.

Erosion Control and Flood Protection

Salt marshes are very effective at minimizing shoreline erosion. Without the extensive root networks of salt marsh vegetation, sediment would quickly be transported away by wind and waves, leading to severe erosion. Salt marsh vegetation also absorbs wave energy and storm surges, temporarily stores floodwaters, and slows river currents. Waterfront homeowners and business owners that have marshes between their property and the ocean benefit from the strong line of defense salt marshes provide against storm damage. Some states are even sponsoring costly salt marsh restoration projects because it is a cost-effective way of protecting coastal communities from storm surges, tropical storms, and Nor'easters.



Water Quality Protection

Streams, rivers, surface runoff, and subsurface groundwater flow all transport pollutants from uplands to marine environments. Salt marshes provide a natural filtration process that can help improve the quality of water that passes through salt marshes before reaching the ocean. Salt marshes take up nutrients such as nitrogen and phosphorus and break down or bind a variety of organic and inorganic pollutants. Although salt marshes are capable of absorbing small levels of excess nutrients, high amounts of certain pollutants such as nitrogen may eventually impair marsh productivity and disrupt food webs. It is important to minimize nutrient inputs to salt marshes in order to maintain valued ecological functions and food webs.

Recreation and Education

Salt marshes offer a wide variety of recreational and educational opportunities for people of all ages. They are popular areas for birding and wildlife viewing. Photographers and artists have long sought the natural beauty of salt marshes for inspiration and solace. Salt marshes are excellent sites for waterfowl hunting and recreational fishing, and kayakers enjoy exploring bays, rivers, and creeks at high tide. As vibrant ecosystems, salt marshes are unique “outdoor classrooms” allow easy access to a rich diversity of plants and animals for natural history and marine biology study and are well suited for coastal ecology lessons.



Threats to New England Salt Marshes

Three hundred fifty years of wetland destruction and pollution have left a lasting legacy on New England salt marshes. Among the challenges facing wetland managers and scientists today are the identification of imperiled salt marshes, the prioritization of sites for restoration, and the development of methods to measure the effectiveness of restoration efforts. The most common threats to salt marshes are changes to natural **hydrology** (including tidal restrictions, ditching, berming, and filling), nonpoint and point source pollution, and coastal development.

Changes to Natural Hydrology: Tidal Restrictions, Ditching, Filling and Stormwater Runoff

Throughout coastal New England, there are vast areas of wetlands that were productive salt marshes until roads or railroads severed their connection to the sea. Transportation routes were built on salt marshes because they were open and flat. Horses and carts were the first to use these routes, followed by steam locomotives in the latter half of the 19th century.

In the 20th century, roadways continued to be built on some of our most valuable wetlands to accommodate automobiles. These roadbeds divided salt marshes into two sections – one with direct unlimited tidal connection to the ocean, and one with restricted or in some cases no access to the ocean (Figure 2.1-B and C). Causeways created for road and railroad crossings are tidal restrictions that have had enormous impacts on landward salt marshes by reducing or eliminating tidal flooding – the force that drives salt marsh ecosystems. Tidal restrictions led to the disruption of natural flooding regimes, alterations to soil and water chemistry, and changes to natural plant and animal communities. These changes led to the establishment and proliferation of invasive species such as the non-native form of *Phragmites australis* (common reed) or *Lythrum salicaria* (purple loosestrife).



Figure 2.1. Man-made impacts such as: A) filling of marsh for development, B) installation of railroad on the marsh surface, C.) tidal restriction, and D) grid ditches effect the Hampton-Seabrook estuary. Photo by SWAMP Inc.

The natural hydrology of salt marshes has also been severely impacted by efforts to manage for them. Colonial settlers first ditched marshes to promote salt hay growth for livestock forage. More recently, resource managers dug deeper ditches to discourage mosquito breeding (Figure 2.1-D). This management action failed because mosquitoes could still find plenty of small, shallow pockets of water on the marsh in which to breed. Ironically, the new ditches drained so much surface water that large areas of marsh habitat could no longer support fish, a primary predator of breeding mosquitoes. By ditching marshes, the amount of permanent standing water

was greatly reduced comprising pool and panne habitat critical to waterfowl, invertebrates, and aquatic vegetation. Ditching salt marshes may also lower the water table and reduce groundwater salinity, increasing the potential for unnatural vegetative changes and the introduction of non-native invasive species on the marsh.

Spoil dumped on salt marshes from activities such as harbor maintenance compact porous salt marsh peat, smother native vegetation, fill in open water habitat and increase marsh elevations.


Environmental Pollution

Human activities release enormous amounts of pollution to the air, water, and soil. The list of pollutants is virtually endless, and their effect on natural ecosystems is not fully understood. The runoff of nutrients, such as nitrogen and phosphorus from fertilizers, septic systems, and farm waste are common nonpoint sources of pollution that in high enough concentrations can change the structure and function of natural ecosystems. Excess nutrients are a particular problem in salt marshes because they eventually lead to **eutrophication**. Industries and combustible engines release numerous heavy metals, such as mercury, lead, and aluminum, which pose lethal and chronic health risks to wildlife and humans. Herbicides and pesticides, which are applied to lawns, gardens, forests, and ponds to kill so-called “nuisance” species, often affect non-target species.

It is beyond the scope of this manual to detail all the types of pollutants and their effects on the environment, but consider this fact: every time it rains, stormwater picks up sediments, nutrients, chemicals, and heavy metals from the landscape and carries these pollutants into storm drains that may lead to streams, rivers, and salt marshes. Salt marshes are depositional areas and therefore are likely to store these pollutants for long periods.

Effects of Human Development

Coastal New England has witnessed unprecedented population growth and urban development over the past three decades. Real estate values have skyrocketed, increasing the pressure on landowners to sell or develop their land. Waterfront property is particularly valuable because of the great views, serenity, and access to the ocean that it can provide. Urban sprawl leads to ever-increasing amounts of impervious surface area, altering surface and groundwater hydrology and increasing nonpoint sources of pollution. The net effect of coastal development and land use change on salt marshes is the loss of upland buffers and



*Coastal development in Hampton infringes on nearby salt marshes.
Photo by NHCP*

new exposure to a wide variety of anthropogenic pollutants and disturbances (Figure 2.1-A). Salt marshes in urban watersheds may receive enormous volumes of stormwater runoff, which can lead to increased erosion, sedimentation, lowered salinity levels, and changes in soil saturation levels. Land use planners project that an additional 25 percent of the New England watershed will be developed by 2025.

The upland buffer and marsh border provide important habitat for a wide variety of wildlife that also utilize the adjacent salt marsh, including many songbird and mammal species. Elimination or alteration of upland buffers indirectly alters bird and wildlife use of salt marshes. Noise pollution and light pollution can affect bird and wildlife behavior. Human disturbance of the landscape may enable potentially damaging species such as *Phragmites australis*, *Lythrum salicaria*, house sparrows, raccoons, and opossums to establish themselves in salt marsh ecosystems and alter natural food webs. Domestic and feral housecats can decimate wild bird populations near residential areas.

Clearly, by virtue of their ecological importance as well as the widespread threats they face, salt marshes are worthy of continued monitoring and research to assess their ecological health, and action to improve or maintain their condition.

Salt Marsh Restoration in New Hampshire

In 1972, Congress passed the Coastal Zone Management Act in recognition of the importance of the nation's coastal resources. The New Hampshire Coastal Program (NHCP) gained federal approval in 1982 and its mission is to “balance the preservation of natural resources of the coast with the social and economic needs of this and succeeding generations.”

Since 1990, NHCP, Ducks Unlimited, the University of New Hampshire, New Hampshire Estuaries Project, additional non-governmental organizations, state and federal agencies, and regional planning commissions have been involved in salt marsh restoration in the nineteen coastal communities. Funding for the NHCP comes primarily from NOAA, administered by the Office of Coastal Resource Management. These projects have restored tidal hydrology by removing tidal restrictions and replacing inadequate culverts, restored marsh elevations by removing fill, restored fish and bird habitat by reestablishing open water habitat, and controlled invasive species such as *Phragmites australis*.

Chapter 3: Elements of Study Design



Parson's Creek, Rye, NH. Photo by NHCP

In recent years, public awareness of the importance and plight of salt marshes has grown. Local residents have become increasingly involved in monitoring programs that support salt marsh preservation and restoration efforts. It can be highly rewarding to take part in salt marsh monitoring – participants can learn about the natural communities of estuarine wetlands and share in public efforts to preserve and protect natural resources. However, it is imperative that volunteers collect data in an organized way so that the information they generate is useful to scientists and resource

managers. This goal is easier than you think! The key to a successful monitoring program is a sound study design. A sound design identifies project goals, and provides specific objectives and methods to achieve those goals. Adherence to procedural protocols ensures data quality. A study design requires that investigators think through and describe how to conduct monitoring to achieve project goals in the form of a document that is read and understood by everybody involved in the monitoring program.

Successful volunteer monitoring programs usually have at least one thing in common: someone to coordinate the monitoring activities, training workshops, logistics, equipment, data sheets, and report preparation. The program coordinator is the hub for the collective effort of the group, and pulls together all the various elements of the project to achieve results and maintain community. The program coordinator usually develops the study design and helps to ensure data quality and consistency, no matter where, when, or by whom the data were collected. Established monitoring programs may be fortunate enough to have funds to compensate the program coordinator, though in many cases the program coordinator is participating as a volunteer.

What To Measure, How, and When

This manual provides guidelines and methods for monitoring three biological parameters (birds, nekton, and vegetation) and four physical/chemical parameters (groundwater, surface water, root-zone salinity, and tidal regime). There are many factors that program coordinators take into consideration when determining which of these parameters to measure. They weigh the pros and cons of each, their relative cost and resources available, and the level of effort and expertise required.

Program coordinators are responsible for selecting parameters, arranging training workshops, and scheduling fieldwork. Volunteer monitors can gain a greater understanding of salt marshes by measuring several parameters. It is better to sample fewer parameters carefully and thoroughly than to sample several parameters at the expense of data quality.

A small amount of good data are far better than a large amount of poor data! Volunteer data are more valuable to resource managers and scientists when collection follows a study design and have used the guidelines and methods provided in this manual.

Site Selection and Sampling Locations

How do resource managers conduct their research? How many sites should be monitored? Should an entire marsh be monitored, or just a portion? How do they decide what areas to monitor? Deciding these factors are important tasks that should be resolved during the development of a study design.

The Comparative Approach

The methods described in this manual are based on use of a comparative approach. There are two primary ways to establish this comparison; the Before-After Comparison and Reference Site-Study Site Comparison (see Box 3.1). When it is feasible, program coordinators will try to incorporate both a Before-After Comparison and Reference Monitoring programs. Combining the two comparative approaches will provide much greater insight into the overall effects of the stressor. The Before-After Comparison allows groups to document the actual response of a marsh to the addition or removal of a **stressor**, and the Reference Site-Study Comparison allows groups to understand restoration targets and provide information and guidance for designing the restoration project.

Reference sites are salt marshes that lack some or all of the disturbances of the study sites or mimic neighboring sites. Reference sites are important because many of the impacts to saltmarshes have occurred over extended periods of time. Many times it is not known what impacted sites were like prior to disturbance. Therefore, reference sites are used as reasonable approximations of conditions in the absence of a particular stressor and are an extremely important part of the study design.

Salt marshes may differ for reasons unrelated to pollution or disturbance. Tidal range, geology, landscape setting, and salinity are just some of the variables that influence salt marsh ecology. Ideally, study sites and reference sites are selected because they are similar in nearly every way except the stressor of interest. Resource managers take into account natural differences between study sites and reference sites and address these differences when analyzing and presenting data.

Example of the Comparative Approach

Example: A group wants to study the effects of a tide restriction on a marsh that will be removed in two or three years. Two different study approaches may be considered.

BEFORE-AFTER COMPARISON

Definition: Study a salt marsh before and after a **stressor** is added or removed.

Application: Study the restricted marsh for one to two years before the removal of the tide restriction and for one to two years afterward. Compare how salt marsh parameters change after the removal. Natural processes respond slowly to change, and restored salt marshes will continue to evolve and respond for a long time; perhaps up to 20 years.

REFERENCE SITE-STUDY SITE COMPARISON

Definition: Compare a salt marsh affected by a particular **stressor** to a similar salt marsh without that stressor.

Application: Use the restricted marsh as the study site and choose a suitable reference site. Usually the unrestricted portion of the salt marsh is a suitable reference site. Compare parameters from the restricted side to the unrestricted side of the salt marsh. This study will provide useful information that will help to plan for the actual restoration and to estimate restoration response. After the removal of the tide restriction, the reference site can serve as a target to help evaluate how the study site is responding.

Four Common Study Areas

Volunteers will be monitoring at four categories of salt marshes: regional reference sites; marshes with tide restrictions; marshes affected by extensive ditching or filling; and marshes affected by pollution, invasive species, and land use.

1. Regional Reference Sites

These salt marshes are generally as pristine as can be found today and include environmental conditions and biological diversity that are representative of a given region. Regional reference sites tend to be large expanses of salt marsh that are owned by conservation entities and are far from development. Ideally, they lack linear or grid ditches that resulted from the Works Project Administration of the 1930s and other ill-conceived mosquito control or drainage projects. Usually, recreational activities such as bird watching, walking, or kayaking are permitted at reference sites, but in general, these marshes experience little human disturbance. Regional reference sites typically represent the best achievable condition for salt marshes in a given region.

2. Salt Marshes With Tide Restrictions

A tide restriction is a reduction in normal tide range resulting from a completely or partially blocked channel. Roads, railroads, and other man-made creek crossings often bisect the marsh into a tidally restricted side and an unrestricted side. The restrictive features of these crossings include undersized or blocked culverts, tide gates, or bridges that restrict full passage of tidal flow. For tide restriction studies, the unrestricted side can be used as the reference site and the restricted side as the study site, because in the absence of the restriction, it is generally assumed that the two sides would resemble each other.



*This 48-inch culvert was providing inadequate tidal flow to over 100 acres of salt marsh at Little River, Rye, NH.
Photo by NHCP*

3. Salt Marshes Affected by Extensive Ditching

Ditching salt marshes drains pool and panne habitat, reducing open water habitat on which many plant, animal, and invertebrate species depend. All of these species can disappear and are part of the natural biological diversity and wealth of a naturally-functioning salt marsh. Ditching the marsh may also lower the water table and reduce salinities, increasing the potential for unnatural vegetative changes and the introduction of non-native invasive species on the marsh.

4. Salt Marshes Affected by Pollution, Invasive Species and Land Use



Meadow Pond, Hampton, NH. Photo by NHCP

Surrounding land uses influence the types and amount of pollutants that enter coastal wetlands, and many researchers are interested in studying the effects of land use and pollution on salt marshes. It is difficult to choose reference marshes for polluted study sites. A common approach is to utilize one or more regional reference sites, with the understanding that there may be some environmental differences between the reference and study sites, such as location in the estuary, soils, topography, or tide exposure. Other reference areas could be parts of the same salt marsh that are farthest from the impacts, or nearby salt marshes whose upland habitat is relatively undisturbed. Project leaders consult with agency scientists or other professionals when selecting reference areas for this category of study sites.

The Evaluation Area

Once study and reference sites are chosen, the next step is to decide where to sample. This task may sound easy for a small salt marsh, but it can be daunting in a 400-acre salt marsh. The study design needs to account for variation in the sizes and environmental condition of reference sites and study sites. To address size variability, scientists have designed an approach to examine comparable portions of reference sites and study sites, called **evaluation areas**. The evaluation area is delineated as follows.

Example: A group is interested in examining the effects of a tide restriction (roadway and culvert) that has bisected a salt marsh into two parts – a six-acre restricted area and a 280-acre unrestricted area. They need to know where to survey the plant community. Here are two common concerns:

1. The unrestricted (reference) area is too big – nearly fifty times larger than the restricted marsh. Sampling the entire 280-acre salt marsh is not feasible or realistic. One plant transect might be a half-mile long!
2. Because of the size difference, you are apprehensive about comparing the restricted marsh to the unrestricted marsh – size alone would likely allow a greater diversity of plants to exist at the reference site.

To address these and other concerns, managers have developed protocols to select representative areas of salt marshes called evaluation areas. Evaluation areas are delineated in a consistent way using specific protocols, and therefore reduce bias associated with size difference between different salt marshes. The location of the evaluation area is also important to isolate and assess the effects of land uses and related impacts like stormwater and fill.

1. From a designated start point on the bank of the salt marsh creek, bay, or salt pond, extend a line along the bank edge for 300 ft (Figure 3.1-1a).
2. At both the start point and the end point, create transects that run perpendicular from the salt marsh banks to the upland edges.
3. The salt marsh and creek habitat that falls between the two transects comprises the evaluation area (Figure 3.1-1b).




Figure 3.1. Delineating evaluation areas. 1a shows the evaluation area on both sides of a tide restriction, and 1b shows the evaluation area along an open-water feature. Courtesy of Mass CZM.

Chapters 4-8 each provide specific instructions on selecting sampling locations for the different parameters.

Cause and Effect

A sound study design should attempt to demonstrate cause and effect. For example, a study design might attempt to answer the question: Is a tide restriction affecting a neighboring salt marsh community? To determine cause and effect, program coordinators must take great care to select suitable reference sites.

Every study has goals, objectives, and limitations. The approach detailed in this manual will indicate if two sites are different, but may not fully explain why they are different. For example, a tide restriction is likely to be a major cause of the reduced biological diversity and increased **abundance** of invasive species, but other factors may be at work as well. There may be a large **groundwater** seep in the study area causing substantial flows of fresh groundwater, which naturally reduces the salinity throughout the system. A population of *Phragmites australis* may have existed at a site for decades, so expectations for removing this invasive species by eliminating the tide restriction may be overly optimistic.



Littorina littorina using creek intertidal habitat. Photo by NOAA

Though we can never be entirely certain of cause and effect in comparative studies, we can overcome some uncertainty by using statistics and weight of evidence. Weight of evidence is the same in ecology as it is in law enforcement -- the more we know about a situation, the more possibilities we can rule out. Volunteer monitoring projects that measure more parameters will be able to build a stronger case for their conclusions. However, volunteer monitoring groups do not often have the time or resources needed to

conduct a study that is intensive enough to build an irrefutable case and as a result, environmental scientists are rarely 100 percent confident about the findings from such studies.

Proposals to carry out restoration activities usually depend on making a strong but not necessarily conclusive case for cause and effect. Although the data may suggest a restoration activity would improve the health/condition or value of a particular marsh, it is impossible to predict the outcome of any specific restoration projects with 100 percent certainty. For this reason, data collection before and after such projects is extremely valuable in improving our understanding of effective restoration strategies. Volunteer monitors can make important contributions to salt marsh protection and restoration without providing academic-level research. In many cases the data provided by volunteer groups help to identify salt marshes that deserve a closer examination, such as a groundwater study, detailed soil and elevation mapping, or further chemical analysis. Another significant function of volunteer monitoring is to track specific parameters like vegetation, fish, and salinity in restoration projects. Observing and documenting the shift from one **community** type to another, or the reduction of invasive species over time, is sometimes as important as understanding exactly why these changes are occurring. Restoration, remediation, protection, and conservation efforts nearly always result from information provided by collaborative efforts between concerned citizens, groups, communities, and professional scientists.

Chapter 4. Data Management, Quality Assurance, and Quality Control

Volunteer monitoring coordinators must ensure that data collected is accurate. Poorly collected samples or data that are carelessly analyzed or presented may mislead people about the quality of the ecosystem. While good data legitimate management decisions, bad data weaken confidence in management actions.

Program coordinators must be able to demonstrate that the data collected by their volunteers are consistent over time, collected and analyzed using standardized and acceptable techniques, and comparable with data collected in other assessments using the same methods. Such assurances include having program coordinators check each incoming data form for decimal errors, missing information, and general problems.

At the start, the program coordinators will institute a strict quality control and quality assurance plan designed to minimize data collection errors, remove data that do not meet rigorous standards, and develop a strategy to present the results. Such a plan will enable program coordinators to stand behind their results and justify their conclusions.

Field data should be recorded on datasheets provided (Appendix A). A separate field data sheet should be used for each site and survey date. Datasheets include information described previously in the protocols (study site, date, station identification, habitat description, species name, total number of individuals, lengths, comments, environmental parameters, etc).

One person should perform all of the data entry so that entries are consistent. If two people are working together, one can observe while the other records information.

When collecting data, there are some things you should keep in mind:

1. Data should be entered neatly, legibly, and thoroughly so that there is not any missing, incomplete, or incorrect information.
2. Make sure that all sections of the datasheet are filled out before moving to the next sampling station.
3. If a variable cannot or was not measured (instrument failure, fish escaped), explain on the datasheet for that section the reason for the missing data. Record the reason/explanation in the field. Do not wait until you return to the office to record the reason or explanation.
4. When the observations have been completed for a site, it helps to review the data sheet to ensure that all the necessary data are accurately recorded and that everything is legible.
5. All field data should be transferred to digital format as soon as possible. Field data are easily incorporated into common spreadsheet or database programs that are designed for comprehensive data management.
6. After the data are entered, it is important to carefully proof the data for errors to insure the data are correct and to maintain quality assurance and quality control of the data.
7. If an error is made in recording data, do not erase. Draw a single pencil line through the incorrect values and enter the correct values.



*Recording data accurately is integral part of monitoring.
Photo by NHCP*

Chapter 5: Ecology of Salt Marsh Vegetation

Plants are the primary food source for salt marsh ecosystems. Most plant material is consumed after it dies as detritus by microbial decomposers and invertebrate consumers. A salt marsh is physically dependent on its plants – plant roots and stems anchor the **substratum** and enable the gradual build up of peat. Plant communities, along with variation in tidal exchange, geology, and chemical parameters such as salinity, shape salt marsh habitat and help determine which species of invertebrates, fish, birds and other animals will be found there.

Salt marshes are an extremely dynamic habitat for plant species because of wide daily and seasonal fluctuations in surface water and root-zone salinity, temperature, and dissolved oxygen. For this reason, the few plants species that are found in salt marshes fill extremely specialized ecological niches. Plant zonation results from species-specific adaptations to physical and chemical conditions.



Seaside Goldenrod
(*Solidago sempervirens*)

Salt Marsh Vegetation Zones

Salt marshes can be extremely difficult places to live because of wide daily fluctuations in salinity, water, temperature, and oxygen. Few plants have evolved adaptations to cope with the extreme conditions of salt marshes. Plant zonation in a salt marsh results from species-specific adaptations to physical and chemical conditions. Looking out on a healthy salt marsh in full summer growth, one can observe distinct zones of plant growth (Figure 5.1). Bands of tall grasses inhabit the saturated banks of creeks and bays, and this zone is bordered by a flat “meadow” of grasses and sedges that may extend landward for a great distance before transitioning into upland habitats where there is a greater diversity of shrubs, flowering plants, and grasses.




Figure 5.1. Plant zonation in northeastern salt marshes. Courtesy of Mass CZM.

Marsh Border: The marsh border is located at the salt marsh’s upland edge and other isolated areas on the marsh where elevations are slightly above the high marsh. The marsh border is usually only flooded at extreme astronomical tides and under irregular conditions such as storm surges or wind-driven tidal inundations, and does not experience waterlogged conditions or severe salt stress. A high diversity of herbs, shrubs, and even trees exists in the marsh border. *Iva frutescens* (high tide bush), *Baccharis halimifolia*

(sea myrtle), *Agropyren pungens* (stiff-leaved quackgrass), *Solidago sempirvirens* (seaside goldenrod), and *Panicum virgatum* (switchgrass) are just some of the many marsh border plants.

High Marsh: The high marsh lies between the low marsh and the marsh's upland border. The high marsh can be very expansive in some areas, sometimes extending hundreds of yards inland from the low marsh area. Soils in the high marsh are mostly saturated, and the high marsh is generally flooded only during higher than average high tides. Plant diversity is low (usually less than 25 species), with the dominant species being the grasses and rushes such as *Spartina patens* (salt hay grass), *Distichlis spicata* (spike grass), *Juncus gerardii* (black grass), and the short form of *S. alterniflora*. Other plant species commonly found in the high marsh are *Aster tenuifolius* (perennial salt marsh aster), and *Limonium nashii* (sea lavender).

Pools: Pannes are located within depressions in the high marsh. They have irregular shallower sloped edges and deeper sections. Pools are generally permanent water holding features in the high marsh that can be vegetated with submerged aquatic species such as *Ruppia maritima* (widgeon grass).

Pannes: Pannes are located interspersed with pools within the high marsh. They are shallower than pools, hold standing water but dry out during extended dry periods. Salinity can reach extremely high concentrations in pannes and only the most salt-tolerant species can exist at panne edges, including *Salicornia spp.* (glassworts), *Plantago maritima* (seaside plantain), and the short form of *Spartina alterniflora*, as well as some blue-green algae.



Salt marsh pool. Photo by NHCP

Low Marsh: The low marsh is located along the seaward edge of the salt marsh. The low marsh is usually flooded at every tide and exposed during low tide. It tends to occur as a narrow band along creeks and ditches, whereas the high marsh is more expansive and is flooded less frequently. The predominant plant species found in the low marsh is the tall form of *Spartina alterniflora* (smooth cordgrass). This species can reach a height of six feet and is very tolerant of daily flooding and exposure.

Why Should Vegetation Be Monitored?

Plant communities respond to human disturbances and subsequent changes to salinity, natural hydrology, invasive species, or pollutant levels. Often, scientists compare a disturbed marsh with an undisturbed marsh to see how vegetation has responded to the disturbance. For example, a study may compare the plant communities in two salt marshes on either side of a railroad bed to try to understand how the restriction altered a previously contiguous ecosystem. Alternatively, a study may record plant communities in one salt marsh over time and watch how vegetation changes in response to the introduction and proliferation of invasive species such as *Phragmites australis* or freshwater wetland species such as *Typha latifolia* (broad-leaf cattail). Many types of disturbance allow plants that could not otherwise live in salt marshes to gain a foothold, reproduce, compete, and perhaps replace native species.



Marsh Orach
(*Atriplex patula*)

When Should Vegetation Be Monitored?

Plants are an important and easy parameter to measure, and sampling should be performed once a season at the time of maximum standing biomass in mid-summer (mid July through August). Sites should be monitored both pre- and post-restoration in years one, two, and every three to five years thereafter.

Where Should Vegetation Be Monitored?

Resource managers and program coordinators will establish evaluation areas to be surveyed. If a creek or river channel bisects the salt marsh (this is the case for most tide restricted sites), volunteers should survey for plants on both sides. In this case, there will be one set of **transects** on each side of the channel (either north and south or east and west).

Transects run from the creek edge to the upland edge, according to a consistent compass bearing (Figure 5.2). For example, all six transects will be laid on a bearing of 275 degrees from the bank to the upland edge.




Figure 5.2 *Vegetation transects at Pickering Brook salt marsh, Greenland, NH. Photo by DU*

How Should Vegetation Be Monitored?

Sampling Methods

❖ Permanent Photo Stations

Photographs taken from permanent stations can provide additional **qualitative** information on changes in a plant community over time. Permanent photo stations should be indicated on maps and marked with permanent field markers. The stations should include views of any restoration activities or structures. Landscape photos are taken at north, south, east, and west compass bearings to cover a panorama of the entire marsh. For stations located at the site of a tidal restriction, bearings that illustrate the downstream marsh are desirable as well. Landscape photographs should include a person or an object for height scale. Photos should be taken at the time of vegetation sampling.

❖ Transect/Plot Method

The general marsh community and species of concern are sampled using $\frac{1}{2}\text{m}^2$ quadrats. In this method, quadrats are placed along the transects at pre-determined intervals.

Equipment

Equipment costs for monitoring vegetation using the transect method are relatively low compared with other parameters. Salt marsh vegetation surveys require a $\frac{1}{2}\text{m}^2$ plot sample quadrat constructed from $\frac{1}{2}$ - inch PVC piping, a comprehensive field guide to wetland plants, a compass, a 50-300 foot tape measure, and flagging. Program coordinators are responsible for building quadrats and supplying all necessary equipment. However, volunteers are welcome to bring their own field guides as well.



Vegetation Survey Protocol

Transect/Plot Method:

1. Please note the section of a transect by placing an N for north, S for south, E for east and W for west after the transect number. *For Example: T1-N = Transect 1, North.* Transects should run north-south or east-west.
2. Starting at the creek edge (the equivalent of 0 feet along the transect), place the quadrat at the following intervals leading away from the creek edge and toward the end of the transect (the upland edge): 0 feet (plot 1), 10 feet (plot 2), 50 feet (plot 3), 100 feet (plot 4), and every 50 feet thereafter until you reach the upland edge. *The first plot is always placed at the beginning of the transect and the last plot is placed at the salt marsh upland border regardless of whether or not the 50 foot interval falls there.*



Examining and recording vegetative cover within a quadrat. Photo by NHCP

3. Facing the upland edge with the tidal channel or creek edge at your back, *always walk on the left side of the tape measure and place the quadrat/plots on the right side.* This ensures that you will not trample plants to be surveyed and that plot position is consistent. Position the quadrats so that the bottom of the left-hand corner of the frame is always located at the designated distance on the measuring tape, such as 0, 10 or 50 ft.

4. Record the transect number, plot number, and distance from the creek edge in the space provided on the data sheet.
5. Record the habitat description within each plot as creek edge, low marsh, high marsh, pool, panne, or upland edge in the space provided on the data sheet.
6. Starting with the first plot on the first transect, identify every plant whose root and stem fall within the $\frac{1}{2}\text{m}^2$ quadrat. Use a field guide to identify plants.
7. Record the abbreviation for the scientific name of each species you have identified in your plot on the data sheet (Table 5.1). Abbreviations consist of two to three letters based on the Latin name of the plant, and can be found on the back of the data sheet.
8. For each species, estimate percent cover by visual examination. Estimate percent bare mud, dead, and areas of open water as well. *The percent cover estimates should always total 100%* (Figure 5.2).
9. If there are species of concern within the plot (Common Reed, Purple Loosestrife, Narrow or Broad-Leaf Cattail), count their stems then measure the height (cm) of the three tallest individuals within each plot.

Cover Type/Species Categories

When collecting vegetation data, please be sure to note the following cover type/species categories:

- ❖ **Bare:** Includes mud and sand. These are areas that are not flooded with water and are devoid of standing live, standing dead, or macroalgae. There can be a thin film of surface water within the bare category.
- ❖ **Dead Vascular Plants:** Identified by species. This category only includes standing dead (attached) plants.
- ❖ **Live Vascular Plants:** (Herbaceous and shrubs) identified by species.
- ❖ **Macroalgae:** Identified as “algae.”
- ❖ **Open Water:** Permanent standing water is identified in plots that are partly within a creek, ditch, pool, or flooded panne.
- ❖ **Wrack:** Wrack is material that has floated into the plot. This is general dead (not attached) plant material.

Plant Identification

Proper identification of wetland plants is an important skill for salt marsh vegetation surveys. Fortunately, New England salt marshes support a low diversity of plants, and the number of species that volunteers will regularly see is limited. Volunteers should try to become familiar with plant **morphology** and plant ecology to become competent with species identification. Several books integrate key information on identification, ecology, and distribution. These books can be invaluable to those without a strong **taxonomy** background.

One of the best publications available for northeastern North America is *A Field Guide to Coastal Wetland Plants of the Northeastern United States* (Ralph Tiner). This field guide provides excellent drawings, clear descriptions, and user-friendly keys for 59 species found in salt and brackish marshes, as well as many other plants found in freshwater wetlands. You may also consult *Life in New Hampshire Salt Marshes: A Quick-Reference Field Guide*, published by the New Hampshire Coastal Program (NHCP) at the Department of Environmental Services (DES).

Always consult the program coordinator if you have trouble identifying a specimen using a field guide. If you are still unable to identify a particular species, you should call the specimen “Unknown Species 1” or “Unknown Species 2” in your field data sheet and place the plant and part of its roots into a re-sealable plastic bag (along with a label) for later identification. Once you have identified the plant, the spreadsheet or database can be adjusted. It may also be a good idea to keep voucher specimens of plants that you are pretty sure you identified properly, but still need some confirmation from someone with more expertise. Vegetation surveys are typically conducted prior to restoration activities, one and two years following restoration activities, and every three to five years thereafter. Several data analysis techniques can be used to detect and analyze changes to a vegetation community over time.

Table 5.1. Common and Latin Names, Habitats, and Abbreviations of Salt Marsh Vegetation


Common Name	Latin Name	Habitat	Code	Common Name	Latin Name	Habitat	Code
Baltic Rush	<i>Juncus balticus</i>	Brackish Areas	JB	Rugosa Rose	<i>Rosa rugosa</i>	Marsh Border	RR
Black Grass	<i>Juncus gerardii</i>	High Marsh	JG	Salt Marsh Aster	<i>Aster tenuifolius</i>	High Marsh	AT
Broad-Leaf Cattail	<i>Typha latifolia</i>	Brackish Areas	TL	Salt Marsh Bulrush	<i>Scirpus maritimus</i>	Brackish Areas	SM
Bushy Knotweed	<i>Polygonum ramosissimum</i>	Marsh Border	PR	Salt Marsh Bulrush	<i>Scirpus robustus</i>	Brackish Areas	SR
Common Glasswort	<i>Salicornia europaea</i>	Pool/Panne	SE	Salt Marsh Gerardia	<i>Agalinia maritima</i>	High Marsh	AM
Common Reed	<i>Phragmites australis</i>	Marsh Border	PA	Salt Meadow Grass	<i>Spartina patens</i>	High Marsh	SP
Creeping Bent Grass	<i>Agrostis stolonifera</i>	High Marsh	AS	Saltwater Cordgrass	<i>Spartina alterniflora</i>	Low Marsh	SA
Glossy Buckthorn	<i>Rhamnus europaea</i>	Marsh Border	RE	Sea Blite	<i>Suaeda linearis</i>	Pool/Panne	SL
Hedge Bindweed	<i>Convolvulus sepium</i>	Marsh Border	CS	Sea Lavender	<i>Limonium nashii</i>	High Marsh	LN
Jewelweed	<i>Impatiens capensis</i>	Brackish Areas	IC	Seashore Alkali Grass	<i>Puccinellia maritima</i>	High Marsh	PUM
Marsh Elder	<i>Iva frutescens</i>	Marsh Border	IF	Seaside Arrow Grass	<i>Triglochin maritima</i>	High Marsh	TM
Marsh Orach	<i>Atriplex patula</i>	High Marsh	AP	Seaside Goldenrod	<i>Solidago sempervirens</i>	High Marsh	SS
Marsh Rush	<i>Juncus effusus</i>	Brackish Areas	JE	Seaside Plantain	<i>Plantago maritima</i>	Pool/Panne	PM
Meadowsweet	<i>Spiraea latifolia</i>	Marsh Border	SPL	Sensitive Fern	<i>Onoclea sensibilis</i>	Brackish Areas	OS
Narrow-Leaf Cattail	<i>Typha angustifolia</i>	Brackish Areas	TA	Silverweed	<i>Potentilla anserina</i>	High Marsh	PAS
Northern Bayberry	Myrica pensylvanica	Marsh Border	MP	Soft-Stem Bulrush	<i>Scirpus validus</i>	Brackish Areas	SV
Olney Three Square	<i>Scirpus olneyi</i>	Brackish Areas	SO	Speckled Alder	<i>Alnus rugosa</i>	Brackish Areas	AR
Poison Ivy	<i>Toxicodendron radicans</i>	Marsh Border	TR	Spike Grass	<i>Distichlis spicata</i>	High Marsh	DS
Prairie Cordgrass	<i>Spartina pectinata</i>	Brackish Areas	SPE	Sweet Gale	<i>Myrica gale</i>	Marsh Border	MG
Purple Loosestrife	<i>Lythrum salicaria</i>	Brackish Areas	LS	Switchgrass	<i>Panicum virgatum</i>	Marsh Border	PV
Red Fescue	<i>Festuca rubra</i>	High Marsh	FR	Unknown Species	<i>Unknown Species</i>		UK
Reed Canary Grass	<i>Phalaris arundinacea</i>	High Marsh	PHA	Widgeon Grass	<i>Ruppia maritima</i>	Pool	RM

- ❖ If macroalgae/microalgae is found, write “algae” in species placeholder of the “Species Present & Percent Cover” column of the datasheet and record percent cover.
- ❖ If the genus of a plant is known but the species is unknown, write the Genus with a “spp” in the species placeholder of the “Species Present & Percent Cover” column datasheet and record percent cover. *For Example: Scirpus spp.*


Figure 5.2. Estimating Percent Cover (%) in Quadrat Plots

SALT MARSH VEGETATION SURVEY
STANDARD COVER CLASSES AND MIDPOINTS FOR ESTIMATING ABUNDANCE


One method for obtaining abundance values is to estimate the percent of a plot occupied by the target plant. To assess percent cover, one estimates the area of the plot frame (1m²) that is covered by all of the leaves, branches, and stems of the target species. Since visual estimates may vary from one person to another, standard cover classes and midpoint abundance values are used to reduce variability. The following figures illustrate nine standard cover classes. For each plot, first identify and list the species present, then for each species determine which figure best describes its cover. Record the midpoint value on the data sheet.




(Trace to 1%)




(11% to 19%)




(46% to 64%)




(2% to 4%)




(20% to 30%)




(65% to 87%)



(5% to 10%)



(31% to 45%)



(88% to 100%)

Chapter 6: Nekton

Salt marshes support diverse and abundant **populations** of creatures that swim. Collectively, these organisms, including fishes and many types of invertebrates, are referred to as **nekton**. This chapter focuses on fish, shrimp and crabs that occupy estuarine wetlands. Salt marshes support most life stages of fish and crabs, which are essential components of the food web and represent a large proportion of the total animal biomass and biological diversity in a marsh. Some species spend only a small portion of their lives in salt marshes, whereas others rarely ever leave. Marsh residents such as mummichogs and four-spine sticklebacks reside in marshes throughout their lives. Transient species use salt marshes during critical development periods such as **spawning** or juvenile rearing and are important seasonal components of salt marsh condition. Transients include forage species such as Atlantic silverside, and commercial and sport species such as winter flounder and striped bass.



Volunteers tally and identify nekton collected using a lift net. Photo by DU

It is challenging to sample nektonic organisms because their distribution and abundance varies greatly throughout the marsh and over time. The use of salt marshes by fish and crabs can vary from tide to tide, marsh to marsh, species to species, and year to year. Even meteorological events such as a full moon or new moon will influence what you are likely to find in a salt marsh. Unlike plants or benthic invertebrates, nektonic animals are highly mobile and difficult to capture. Despite these challenges, fish and crabs are fun to study and learn about, can be important indicators of salt marsh condition, and in many cases are the impetus for marsh restoration.

The goals and objectives of a study will dictate the sampling methods used. Monitoring efforts will usually attempt to gather baseline information on species presence and relative abundance to evaluate potential differences between reference and study sites and allow evaluation of monitoring techniques. Volunteers can easily obtain qualitative information about common marsh species. **Quantitative** estimates are possible as volunteers gain experience with salt marsh sampling.

Why Should Nekton Be Monitored?

Scientists do not fully understand the influence of marsh degradation on fish and crabs, though they continue to investigate this important topic. Tide restrictions may alter fish and crab communities by reducing habitat availability, accessibility, and quality on the restricted side. Many species are **sensitive** to changes in dissolved oxygen, salinity, and nutrient levels that result from pollution and surface runoff. Changes to salt marsh vegetation resulting from upland human disturbance, alterations to natural hydrology, or invasive species may affect fish or crabs that require native or natural plant communities.

When Should Nekton Be Monitored?

Most study sites will have one to two daytime survey events per year; one in early summer and another in late summer-early fall unless there are species or processes unique to other seasons that are of interest. Fish and crabs are typically sampled from June through September, although surveys can be conducted year round. Nekton sampling in ditches should occur at the same relative tidal stage or tidal cycle and should occur only after the marsh surface is empty of tidal water. Sites should be monitored both pre- and post-restoration in years one, two, and every three to five years thereafter.

Where Should Nekton Be Monitored?

The number of individual stations required depends on the habitat under examination, but volunteers will probably sample a total of 20 stations during each survey event (10 stations at pool/panne habitats, and another 10 at creek/ditch habitats). Once sampling stations have been chosen, their locations should be marked with stakes or flagging. Their locations can also be recorded using high-accuracy GPS.


How Should Nekton Be Monitored?

Sampling Methods

- ❖ **Ditch Nets** should be used to sample nekton in small creeks and man-made ditches.
- ❖ **Lift Nets** should be used to sample nekton in pools, pannes and plugged ditches.

Equipment

There are a variety of equipment and methods used to collect salt marsh nekton, each suitable for different conditions, habitats, and target organisms. You will need aerial photos of study sites showing boundaries of study areas and approximate locations of creeks/ditches and pools/pannes, or program coordinators will mark their locations on an aerial map. You will also need “ditch nets”, “lift nets”, small rulers, containers to hold the fish, meter sticks to measure water in the nets, small “dip” nets, and a water quality meter.



Nekton Survey Protocol

Ditch Nets

There are two “doors” on the open ends of the net, which when pulled, rise up to close off the ends of the net, enclosing an area of water that is approximately one linear meter long and as wide as the ditch.

1. Place ditch net at the station locations at least 30 minutes before sampling. This is done to minimize any disturbance to nekton caused by placing the net in the ditch. This usually means that the nets are placed prior to the incoming high tide, so that they can be pulled at the incoming or outgoing high tide.
2. Setting up a ditch net usually requires two people, each standing on opposite sides of the ditch or pool. One person will take stakes labeled “A” and “B” and place the stakes into the bottom of the ditch close to the side of the ditch. The other person will take stakes labeled “C” and “D” and place them on the opposite side of the ditch. The net should be stretched tight between stakes “A” and “B” and stakes “C” and “D” so that approximately 1 meter length of ditch is sampled (Figure 6.1).




Figure 6.1. *Measuring dimensions of a ditch net.*
Photo by DU

3. Pull the ripcords to make sure that the lines are not tangled and that the doors will pull up smoothly and quickly.
4. Push the doors and the center of the net down into the bottom of the ditch with the meter stick. Make sure that the net lay down on the bottom of the ditch, so that fish passage through the net is not impeded.
5. Measure the distance between all the stakes (*e.g.* “A” to “B”, “B” to “C”, “C” to “D”, and “D” to “A” and the diagonal distance between stakes “A” and “C”) and record these on the datasheet (Appendix C). These distances are measured when the ditch net is placed in the ditch and are necessary to calculate the area of water that is sampled. Measure these distances at approximately $\frac{3}{4}$'s of the height of the ditch net (or where $\frac{1}{3}$ - $\frac{3}{4}$ of tidal water will be in the net when pulled).
6. Lay the ripcords straight away from the ditch net along the marsh surface as far from the net as possible without pulling on the doors.
7. Record the time that the net was installed on the data sheet, in military time.
8. Two people are required to pull a ditch net. At the incoming or outgoing high tide (when approximately $\frac{1}{3}$ - $\frac{3}{4}$ of the net is filled with water, and after at least 30 minutes from the time it was installed), pull the ditch net.



Approaching a ditch net after drawing the ends closed. Photo by DU

Quietly and slowly approach the net from opposite sides of the ditch, with one person on either side. Upon reaching the ripcords, kneel and wait quietly for approximately two minutes. Do not handle the ripcords during this time, as the vibrations on the cords can be transmitted to the stakes and possibly disturb nekton that are in the net. At a pre-determined signal, both people quickly pull on the ripcords. The doors of the net will pull up, enclosing nekton within the ditch net. Once the edges have been fully raised, both people should approach the net. Keep the ripcords taught while approaching the net, so as to prevent any escape.

9. Quickly lift the ditch net out of the ditch and onto the marsh surface. The best way to do this is to have both people pull the stakes out simultaneously (while still maintaining pressure on the ripcords). One person should hand two stakes to the other so that person can lift the net out of the ditch and onto the marsh surface. It is important to quickly pull the stakes and net out of the ditch. This creates a bag of netting in the center of the net where nekton are trapped (because otherwise, the nekton could still leave through the sides of the doors).
10. In military time, record the time that the ditch net was collected on the data sheet.
11. Lay the net out on the marsh surface and collect and place all nekton captured into a container of water. Separate crabs and other potential predators by placing them in another container.
12. For each sample, measure (to the nearest mm) up to 15 individuals of every species. Measure fish from the tip of the snout to the tip of the caudal fin; shrimp from the tip of the rostrum to the tip of the telson; and crabs from the distance between the two furthest points across the carapace. Count and measure dominant fish species such as mummichogs (*Fundulus heteroclitus*) in two age classes: juveniles (<45 mm) and adult male and female (>45 mm). Enter juveniles and adults separately in the spaces provided on the data sheet.
13. Place those individuals measured and counted into a separate container of water. When finished measuring and counting, return the nekton to the water as soon as possible to limit mortality.
14. Collect the depth (cm) of the water in the tidal creek or ditch, just after collecting the net, and record in the space provided on the data sheet. Depth is measured to calculate the volume of water that is sampled, and ultimately to calculate fish per m³ at each station.
15. During nekton collection, collect water quality information, using a YSI instrument, and determine water temperature (degrees Celsius), salinity (parts per thousand), dissolved oxygen (milligrams per liter and percent saturation), and record in the spaces provided on the data sheet.



*Juvenile mummichog; 37 mm in length.
Photo by DU*

Lift Nets

1. Select a section of a pool/panne for sampling so that the lift net may be placed on the bottom and pulled easily with minimal disruption to water and sediment.
2. Place lift net into position at least 30 minutes prior to collecting.
3. Two people are required to set a lift net. Unwrap net so that weighted cords are on the bottom, facing the ground. Make sure there are no holes, tears, or bunches in the net.

4. With one person on each short end, drape the net on the surface of the water. Weights should slowly lower the net to the bottom.

5. Starting at the center, gently press the entire width of the net into the mud using a wooden stake securing the net to the mud bottom. Secure the rest of the net by pressing it into the mud at one to two foot intervals down the length. (Tip: At one to two feet from center, place the wooden stake gently on the net. Move net slowly toward the center and depress into mud. This will ensure that the center does not shift out of the mud).



*An installed lift net with well placed drawstrings.
Photo by DU*

6. Make sure that all edges and corners are lying flat or depressed into the mud. Pockets will leave areas for fish to hide underneath.

7. Lay the ripcords out on the marsh surface as far from the net as possible without pulling on the net.

8. Measure the length and width of the net (cm) and record in the spaces provided on the data sheet.

9. Measure the depth of the pool (cm) at the most common depth and record in the space provided on the data sheet.

10. In military time, record the time the lift net was installed on the data sheet.

11. Two people are required to pull a lift net. Both individuals should approach ripcords quietly and slowly to reduce disturbance to the pool and net. Once at the ripcords, kneel and wait approximately two minutes. *Do not touch cords until signal is given. Vibrations will cause fish to retreat from net.* At a pre-determined signal, pick up the rope and



quickly pull up while stepping backwards to lift the net out of water. After two to three seconds, most of the water will have drained from the net and the narrow width may be folded to form an “envelope” to prevent fish from escaping.

12. Keeping net folded, bring the net to a stable high marsh surface.

13. Lay the net out on the marsh surface and collect and place all nekton captured into a container of water. Separate crabs and other potential predators by placing them in another container.

14. For each sample, measure (to the nearest mm) up to 15 individuals of every species. Measure fish from the tip of the snout to the tip of the caudal fin; shrimp from the tip of the rostrum to the tip of the telson; and crabs from the distance



Counting and measuring fish. Photo by NHCP.

between the two furthest points across the carapace. Count and measure dominant fish species such as mummichogs (*Fundulus heteroclitus*) in two age classes: juveniles (<45 mm) and adult male and female (>45 mm). Enter juveniles and adults separately in the spaces provided on the data sheet.

15. Place those individuals measured and counted into a separate container of water. When finished measuring and counting, return the nekton to the water as soon as possible to limit mortality.
16. During nekton collection, collect water quality information, using a YSI instrument, and determine water temperature (degrees Celsius), salinity (parts per thousand), dissolved oxygen (milligrams per liter and percent saturation), and record in the spaces provided on the data sheet.

Table 6.1. Common Salt Marsh Fishes and Crabs

Common Name	Scientific Name	RES	TRA	FRE	BRA	MAR	ANA	CAT
American Shad	<i>Alosa sapidissima</i>		X	X	X	X	X	
Atlantic Herring	<i>Clupea harengus</i>		X	X	X	X		
Atlantic Silverside	<i>Menidia menidia</i>		X					
Fourspine Stickleback	<i>Apeltes quadracus</i>	X		X	X	X		
Green Crab	<i>Carcinus maenas</i>	X			X	X		
Mummichog	<i>Fundulus heteroclitus</i>	X		X	X	X		
Striped Killifish	<i>Fundulus majalis</i>	X		X	X	X		
Ninespine Stickleback	<i>Pungitius pungitius</i>		X		X	X		
Threespine Stickleback	<i>Gasterosteus aculeatus</i>		X	X	X	X		
American Eel	<i>Anguilla rostrata</i>							X
Sheepshead Minnow	<i>Cyprinodon variegatus</i>	X						

Abbreviations:

RES = Residential

TRA = Transient

FRE = Freshwater

BRA = Brackish

MAR = Marine

ANA = Anadromous

CAT = Catadromous

Chapter 7: Birds

Birds are the most conspicuous animals inhabiting New England salt marshes because they fly around, sing, and attract attention. For centuries, scientists and naturalists have studied avian life history, behavior, environmental requirements, and responses to environmental disturbance and pollution. The concept of using birds as sensitive environmental indicators has long been established: miners brought caged birds into mines to serve as indicators of air quality, giving rise to the expression “canary in a coal mine.” More recently, scientists have studied how nesting, hatching, and fledging success can reflect environmental conditions. This chapter provides the guidelines and methods needed to conduct a monitoring for salt marsh birds, and discusses how birds may be used as environmental indicators.



Parker River National Wildlife Refuge, Mass. Photo by USFWS

Why Should Birds Be Monitored?

Wetland birds require certain types of habitats for different aspects of their lives such as nesting, feeding, or migration. Salt marshes offer a variety of habitats such as mudflats, open water, and various types of vegetation. Birds have evolved a variety of adaptations to exploit the resources in these habitats. For instance, herons and egrets have long legs well suited for wading in shallow water and beaks suited for catching fish and invertebrate prey. Habitat diversity in salt marshes results from a variety of physical, chemical, and biological variables. Alterations to physical variables such as hydrology, chemical variables such as salinity, or biological variables such as vegetation will affect the type and distribution of habitats in a salt marsh, and therefore the biological communities that can live there.

Humans may alter the habitat that a bird requires. For instance, salt marsh sharp-tailed sparrows (*Ammodramus c. caudacutus*) require suitable densities of *Spartina patens* and *Spartina alterniflora* for nesting and feeding. As discussed in previous chapters, alterations to natural hydrology or salinity regimes may reduce the availability of these vegetation types. Humans may also alter the abundance of important prey items. For instance, herons and egrets require high densities of fish, and excessive pollution or alterations to natural hydrology might reduce fish populations to the point where herons and egrets cannot get adequate nutrition. Birds that require specific habitats or conditions -- called **specialists** (egrets and sandpipers) -- may avoid salt marshes that have been altered by disturbance or pollution, while birds that can tolerate a wide range of conditions -- called **generalists** (gulls and crows) may thrive in these areas.

Birds are long-lived and highly mobile, and over the course of their lives can learn to recognize favorable locations for breeding, nesting, feeding, and migration. Scientific research has shown that birds will choose wetlands that have the best conditions to meet their needs. Birds are usually able to recognize if the vegetation is suitable for nesting, or if prey abundance is sufficient for themselves and their young, and if predation risk is low. Birds prefer to avoid stressful situations, and will usually only occupy unfavorable

habitats if competition forces them to leave favorable areas. Human land use may affect a bird's decision to inhabit a particular salt marsh - - some birds avoid wetlands near highways and urban areas because of noise pollution, and large numbers of cats from suburban developments might be a deterrent for birds. Birds that seek out favorable habitats will be more healthy, more likely to have healthy offspring, and may be at lesser risk of mortality.

Birds are just one of several types of organisms that inhabit salt marshes. Although there are shortcomings to using birds to assess the condition of salt marshes, they can yield insight that may be overlooked by studying only plants, invertebrates or fish.

Bird Identification

Bird identification requires careful visual observations and keen auditory skills. Although some birds are very distinctive, many others look similar and often confuse even the most skilled observers. Some



Egrets and greater yellowlegs in salt marsh pool.
Photo by DU

difficulties may arise from the fact that many birds molt twice a year and the appearance of their plumage changes. Juvenile birds that have not developed adult characteristics will often be difficult to identify. Recognizing birdcalls can be an important means of identifying species, (particularly cryptic species), but commonly available recordings often do not include all species you will encounter. Most birds that regularly use the salt marsh, with the exception of the sparrows, are generally silent until disturbed. Many bird identification books are available, and rely on a suite of illustrations, photographs, and descriptions. For best results, gather the bird identification materials you are most comfortable with and use a variety of clues (shape, posture, size, coloration, behavior, habitat, and birdcalls) to identify species or groups of closely related species.

As stated previously, actual bird surveys require constant attention by the observers so that they do not overlook

any species or miscount individuals. Volunteers should spend several days practicing in the field before actually conducting field surveys to familiarize themselves with the birds and survey conditions. During these “practice runs,” volunteers should follow a series of steps to narrow the range of possibilities for any given species and arrive at the proper identification; these steps are outlined below. Volunteers will not conduct actual field surveys until they are proficient with identifying birds by sight.

The first step for visual observation is to determine what general type of bird you are looking at. You should know key characteristics for a few basic groups of birds, based largely on shape and posture. Familiarize yourself with the main group of birds, many of which you probably already know to some extent, so that you can ask yourself simple questions such as:

- ❖ Is it gull-like? (gulls, terns)
- ❖ Is it duck-like? (ducks, geese, swans)
- ❖ Is it hawk-like? (Ospreys, eagles, hawks)
- ❖ Is it a wading bird with long legs? (herons, egrets)
- ❖ Does it run along the ground like a sandpiper? (“shorebirds” - sandpipers, plovers)
- ❖ Is it a perching bird? (Large group of birds, which includes mostly songbirds)

Once you identify the general group a bird belongs to, consult a bird book to find the species that the bird most resembles. You should be familiar with the organization of your book so you can quickly reach the appropriate section and spend more time comparing closely related species. It is sometimes helpful to take notes or photographs to assist in identification. Using common species for comparison, you should focus on details such as size (bigger or smaller than a robin?), shape (long and thin or plump and round?), coloration (brown? what shade?), and any distinguishing marks or features (any streaking or other noticeable marks?).



Black ducks and mallards at the marsh shoreline. Photo by DU

Pay attention to the bird's behavior, including feeding, roosting, flying, and the types of habitat it occupies. Be mindful of the time of year, because in the summer and fall you are likely to see immature birds or post-breeding adults whose plumage is different than what is illustrated in most books. Some groups of birds are difficult to identify, including sparrows, flycatchers, young gulls, fall warblers, and starlings (due to the many variations in their plumage).

Birdcalls are frequently more difficult to learn than visual cues, but knowing the calls will dramatically increase your ability to identify birds in the field. This is particularly true for birds that are cryptic or otherwise difficult to see because of weather, darkness, or heavy cover. Listening to birds as they are calling is perhaps the best way to learn their calls, because this “hands on” approach will enable you to create strong and long-lasting associations between a species’ appearance and call. But remember that most birds that regularly use the marsh are silent unless disturbed.

Bird Activity and Habitat

You can determine the importance of a wetland to a particular species by recording its activity and habitat usage. Here are some examples:

- ❖ Some birds (such as gulls, ducks, and hawks) will fly over a wetland on their way to somewhere else. You should not count birds that are cruising high above a wetland unless it looks like they are hunting (such as an osprey looking for prey).
- ❖ Some birds (such as swallows, swifts, and flycatchers) cruise at low altitude over the marsh and feed on aerial insects. You should count these low-flying birds.
- ❖ Some birds feed almost entirely within the salt marsh yet nest and perch in adjacent wetland buffers because the vegetation may be more dense and protected. You should count birds in the marsh and the upland buffer and record where you observed them.
- ❖ Do your best not to count the same individual twice. Since birds may be flying around, this may be difficult. If a bird flies off in one direction and out of sight and then another individual of the same species appears to fly back from the same area, it is possibly the same bird. Use your best judgment.
- ❖ If you cannot identify a bird, do not spend too much time looking it up in your book, since you will miss other birds that fly by. Jot a few notes and try to figure it out later – such as color, general size and bird group – gull, waterfowl, shorebird, etc.

When Should Birds Be Monitored?

Recent studies in Maine found a strong correlation between the tide cycle and sharp-tailed sparrow singing activity. Sharp-tailed sparrows sing much more just after the new moon tide. Therefore, observers should focus survey activity around these periods to ensure that each point is sampled during this time period.

How Should Birds Be Monitored?

Sampling Methods

Scientists use two types of field protocols to monitor birds: walking transects and point counts. Walking transects and point counts have been used in all New England states, New Brunswick, and Nova Scotia in order to ensure consistency and comparability in data collection. This data will provide the baseline estimates for salt marsh breeding bird distribution and abundance and aid in establishing a long-term monitoring program for these under sampled species. Therefore, it is important to accurately record where each point is located in a marsh so comparisons with future surveys will be valid.

Point Counts

Surveys are conducted from a single vantage point overlooking the marsh, and require observers to record all species and individuals seen or heard within a fixed amount of time.

1. Salt marsh bird surveys are created to adequately cover the marsh's study area. In large marshes a line of sampling points may be on the outside edge of the marsh and then move further into the marsh as the route continues, forming a loop so that the observer ends up at the starting point without having to backtrack.




Figure 7.1. Diagram of bird observation points.

2. The points will be at least 25m from the upland edge, which is unsuitable or non-marsh habitat, and will include a 50m radius of observation. Point centers should be 150m apart, leaving the outside edge of each point 50m apart (see Figure 7.1).
3. This creates a sampling design where each point is independent from another, thus avoiding counting the same individuals twice. In small marshes or fringing marshes, there may be sampling in unsuitable habitat or close to an edge.
4. Some of the sampling points will have a paired flag to signify the 50m edge to assist with estimating distance to birds.


Walking Transect

These are sometimes used alone or in conjunction with point counts. Birds are counted along a standardized path between point counts.

Equipment

Of the parameters covered in this manual, birds are among the least costly to monitor. Observers only need 5 pieces of equipment to monitor birds: binoculars, field guides, pencils, a clipboard and patience! Binoculars range in price, but most backyard birdwatchers and outdoor enthusiasts already have a pair. Recommended bird identification field guides include:

- ❖ *Field Guide to the Birds of North America*
- ❖ *The Sibley Field Guide to Birds of Eastern North America*
- ❖ *Peterson's Guide to Birds*
- ❖ *Field Guide to Birds East of the Rockies.*



Bird Survey Protocols

Point Counts/Walking Transect Combination

1. Point counts should be conducted from sunrise to 11 a.m. during the breeding season, and should occur at various parts of the day during the non-breeding season surrounding the high tide. If a survey is conducted late at one session, be sure to conduct the next survey earlier in the day. Point observations should be conducted for 10 minutes each. Points should not be run in high winds or strong rain, but a light drizzle is acceptable.
2. Field surveys require constant attention, and there is little time for looking at books during fieldwork. Identify, count, and record on the data sheet all birds observed during the 10-minute point. (Table 7.1)
3. Record when the bird was observed during the 10 minutes; between zero to three minutes, three to five minutes and five-10 minutes. (This allows researchers to compare these data to Breeding Bird Survey data and helps researchers develop a long term monitoring program). It is important to note that when the count is divided into time intervals, only count the new birds detected in each time interval (i.e., once an individual bird is recorded, do not count it again). So if you observed a Willet in the first 30 seconds and then detected the SAME bird 5 minutes later, that is still one Willet. *Do not count that bird again in the five -10 minute time period.*


- Use the forestry dot method to count individuals, then at the end of the day, go through the data sheets and add up the dots for each cell.

Each point and line counts as an individual bird of the same species. When counting, fill the corners in first, then connect them with lines until there are 10 symbols. This avoids many mistakes made using the slash and cross method of tallying. For example:



Record whether the individual was within one of three distance categories (0-50m; 50-100m, or beyond 100m). Most of the analyses will be conducted on species within 50m, but researchers also want to be able to include **vagile** species and species with large territories.

NOTE: Salt marsh sparrows are a species that rely heavily on salt marshes in New England. Until recently, little information was available on their habitat requirements. For this reason, if a salt marsh sparrow is heard singing, please indicate this on the data sheet, and circle the dots that represent birds that were seen only (did not vocalize). This gives researchers the ability to pull males only from the data. All non-circled dots indicate that the individual was heard or both seen and heard. This will not be used in the overall monitoring design but may be a useful guide for future research.



Data Entry in the Field

One person should perform all of the data entry so that entries are consistent. If two people are working together, one can observe while the other person records information.



*Volunteer monitors observing birds.
Photo by DU*

Table 7.1. Birds Observed Using Salt Marshes in New Hampshire

Species	AOU Code	Species	AOU Code
American Black Duck (<i>Anas rubripes</i>)	ABDU	Killdeer (<i>Charadrius vociferous</i>)	KILL
American Crow (<i>Corvus brachyrhynchos</i>)	AMCR	Herring Gull (<i>Larus argentatus</i>)	HEGU
American Goldfinch (<i>Carduelis tristis</i>)	AMGO	King Rail (<i>Rallus elegans</i>)	KIRA
American Robin (<i>Turdus migratorius</i>)	AMRO	Least Bittern (<i>Ixobrychus exilis</i>)	LEBI
American Tree Sparrow (<i>Spizella arborea</i>)	ATSP	Least Sandpiper (<i>Calidris minutilla</i>)	LESA
Bank Swallow (<i>Riparia riparia</i>)	BANS	Least Tern (<i>Sterna albifrons</i>)	LETE
Barn Swallow (<i>Hirundo rustica</i>)	BARS	Lesser Yellowlegs (<i>Totanus flavipes</i>)	LEYE
Belted Kingfisher (<i>Ceryle alcyon</i>)	BEKI	Mallard (<i>Anas platyrhynchos</i>)	MALL
Black-capped Chickadee (<i>Parus atricapillus</i>)	BCCH	Marsh Wren (<i>Cisthorus palustris</i>)	MAWR
Black-crowned Night Heron (<i>Nycticorax nycticorax</i>)	BCNH	Mourning Dove (<i>Zenaida macroura</i>)	MODO
Blue Jay (<i>Cyanocitta cristata</i>)	BLJA	Nelson's Sharp-tailed Sparrow (<i>Ammodramus n. subvirgatus</i>)	NSTS
Blue-wing Teal (<i>Anas discors</i>)	BWTL	Northern Harrier (<i>Circus canesus</i>)	NOHA
Bobolink (<i>Dolichonyx oryzivorus</i>)	BOBO	Osprey (<i>Pandion haliaetus</i>)	OSPR
Brown-headed Cowbird (<i>Molothrus ater</i>)	BHCO	Purple Martin (<i>Progne subis</i>)	PUMA
Canada Goose (<i>Branta canadensis</i>)	CAGO	Red-Winged Blackbird (<i>Agelaius phoeniceus</i>)	RWBL
Chimney Swift (<i>Chaetura pelagica</i>)	CHSW	Ring-billed Gull (<i>Larus delawarensis</i>)	RBGU
Clapper Rail (<i>Rallus longirostris</i>)	CLRA	Saltmarsh Sharp-tailed Sparrow (<i>Ammodramus c. caudacutus</i>)	SSTS
Common Grackle (<i>Quiscalus quiscula</i>)	COGR	Savannah Sparrow (<i>Passerculus sandwichensis</i>)	SAVS
Common Tern (<i>Sterna hirundo</i>)	COTE	Seaside Sparrow (<i>Ammodramus maritima</i>)	SESP
Common Yellow Throat (<i>Geothlypis trichas</i>)	COYE	Semipalmated Sandpiper (<i>Calidris pusilla</i>)	SESA
Cooper's Hawk (<i>Accipiter cooperii</i>)	COHA	Short-billed Dowitcher (<i>Limnodromus griseus</i>)	SBDO
Double-crested Cormorant (<i>Phalacrocorax auritus</i>)	DCCO	Snowy Egret (<i>Egretta thula</i>)	SNEG
Eastern Bluebird (<i>Sialia sialis</i>)	EABL	Song Sparrow (<i>Melospiza melodia</i>)	SOSP
Eastern Kingbird (<i>Tyrannus tyrannus</i>)	EAKI	Spotted Sandpiper (<i>Actitis macularia</i>)	SPSA
European Starling (<i>Sturnus vulgaris</i>)	EUST	Swamp Sparrow (<i>Melospiza georgiana</i>)	SWSP
Glossy Ibis (<i>Plegadis falcinellus</i>)	GLIB	Tree Swallow (<i>Tachycinets bicolor</i>)	TRES
Greater Yellowlegs (<i>Totanus melanoleucus</i>)	GRYE	Virginia Rail (<i>Rallus limicola</i>)	VIRA
Great Blue Heron (<i>Ardea herodias</i>)	GBHE	Willet (<i>Catoptrophorus semipalmatus</i>)	WILL
Great Egret (<i>Ardea alba</i>)	GREG	Willow Flycatcher (<i>Empidonax traillii</i>)	WIFL
Great Black-backed Gull (<i>Larus marinus</i>)	GBBG	Yellow-rumped Warbler (<i>Dendroica coronata</i>)	MYWA
Green-wing Teal (<i>Anas carolinensis</i>)	GRWT		
Green Heron (<i>Butorides striatus</i>)	GRHE		

For additional bird codes refer to <http://www.pwrc.usgs.gov/BBL/manual/sname.htm>

Chapter 8: Soil Characteristics

There are several types of water chemistry studies, including investigations of **ambient water quality** trends that typically involve several parameters, or specific investigations of suspected sources of pollution or parameters of interest. Ambient water quality sampling is usually conducted concurrently with the nekton monitoring methods described in this handbook, and the parameters of interest include salinity, dissolved oxygen, pH, and temperature. This chapter reviews methods for measuring two soil characteristics: soil salinity and sediment elevation.



*Volunteers monitor the salinity of well samples.
Photo by NHCP*

Soil Salinity

Tidal inundation and freshwater sources control **salinity regimes** in salt marshes. Salinity is highest in areas of estuaries closest to the ocean and in pools or pannes within salt marshes, and gradually declines in a landward direction as the effects of tidal inundation diminish. Rainfall is another source of verification and can reduce salinity in pools substantially. Perhaps the most recognizable consequence of salinity regimes in salt marshes is the vegetation zonation patterns (see Chapter 4).


Measurements of salinity can help to explain the diversity, distribution, and abundance of plants and animals in a salt marsh. Salinity is also a critical parameter to measure when investigating any type of tidal restriction (or man-made alteration to marsh hydrology). Careful measurements before and after the removal of a tide restriction can provide an excellent indication of the success of restoration efforts. Most often, the primary goals of programs that seek to restore tidal flow are the reestablishment of natural salinity regimes for the restoration of communities.

When Should Salinity Be Monitored?

Soil salinities should be obtained throughout the entire area to be restored and at an appropriate reference marsh. Sampling should be performed at low tide every three weeks between the beginning of the growing season (April or May) to late-season (September or October) including both spring and neap tides. Soil salinity monitoring may continue at post-restoration sites for several years until results of the restoration activity can be determined. If annual assessments show positive results from the restoration, it can be omitted in some years (for example: pre-restoration, year one, two, four, five, and seven post-restoration).

Where Should Salinity Be Sampled?

A simple layout of at least five stations per marsh unit (i.e. upstream and downstream of tidal restriction or scattered) will be established as a minimum for sampling soil salinity. Along the axis of the main channel one station would be placed close to the restriction, one near the predicted head of tide (after restoration) and one equidistant between the two. These three stations would be located in high marsh approximately 3m to 4m from the tidal channel. Two more stations would be placed between the central station and the upland in high marsh. Soil water should be collected from 5 to 20cm depths (0 to 5cm samples are not practical except with soil cores).




*Example of a salinity monitoring layout
X's mark salinity wells.*

How Should Salinity Be Monitored?


Equipment

Soil salinity is a relatively inexpensive parameter to incorporate into a monitoring program. Wells to obtain soil salinity are constructed from 19mm diameter CPVC plastic pipe with seven pairs of 4mm holes at sediment depths between 5cm to 20cm. The base of the 35cm pipe is sealed and the top is capped with two right angles in sequence. This prevents rain or floodwaters from entering the well while maintaining ambient air pressure.



Sampling Method

1. Remove the elbow-shaped cap from the well and insert the syringe tubing to the base of the well.
2. Lift the tubing about an inch away from the base to minimize the presence of mud and silt in the sample.
3. Extract a sample from the well by pulling the circular knob at the top of the syringe and drawing as much water into the syringe as possible.
4. Shake the syringe to reduce stratification (layering) in the water.
5. Open the daylight sample plate. Place the end of the tubing onto the refractometer prism and gently push the plunger until a few drops of water fall onto the refractometer.
6. Close the daylight plate so the plate comes into contact with the prism surface. The sample should spread completely over the prism surface. Air bubbles or an insufficient sample will be hard to read. If this occurs, repeat the procedure, applying more of the liquid sample.
7. Hold the refractometer by the rubber grip and point the prism end of the refractometer to a light source and observe the field of view through the eyepiece. Focus the eyepiece by turning the cross-striped portion of the rubber eyepiece guard either clockwise or counter-clockwise until the scale becomes clearly visible.



8. A horizontal boundary line separating the blue field of view (top) and white field of view (bottom) will appear in the field of vision. Adjust the angle of unit to the light source until line is sharp and distinct. The values on the scale are specific gravity and parts per thousand (ppt).
9. Measure the sample to the nearest parts per thousand (ppt). Record the information in the space provided on the data sheet for “sample 1”.
10. Wipe the refractometer prism surface dry and repeat steps 5 and 6 two more times for a total of three salinity readings.
11. Do not allow saltwater to remain anywhere on the unit. When through, wipe clean with a cloth dampened with fresh water then wipe dry.

Sediment Elevation

Measuring the surface elevation of the marsh may assess net balances in critical soil processes that allow salt marshes to persist over time. Loss in elevation indicates peat degradation, whereas gains may be due to accretion at the surface or peat development below the surface. Standard survey techniques are unable to measure short-term changes in the sediment elevation (two to three years), but are adequate for documenting long-term change (10 years or greater). Short-term changes in the sediment elevation of salt marshes around the world are being monitored using sediment elevation tables (SET). Installation of the SETs is difficult and requires professionals. In New England, more than 30 stations exist in several salt marshes. Data are collected to provide baseline information, assess projects to restore hydrology, and assess sea level rise.

Chapter 9: Hydrology

The presence, type, and potential hydrologic effects of tide restrictions are critical information for salt marsh monitoring and assessment. Tidal influence is an important parameter to measure, and along with salinity can provide a very good understanding of the effect of tide restrictions on the physical and chemical nature of salt marshes.

Tidal Flow

As discussed earlier, tide restrictions usually result from the construction of a travel route over a salt marsh, particularly where a bridge or culvert is installed on the tidal creek. Tidal crossings are restrictive if they block or inhibit water from flowing freely from one side of the marsh to the other, resulting in a reduction of tidal influence on the landward, or restricted side of the estuary. The seaward, or unrestricted, side of the estuary is a good indication of what the restricted side would resemble in the absence of the tide restriction. In tidal influence studies, the unrestricted marsh is usually the reference marsh and the restricted marsh is usually the study marsh. A comparison of tidal ranges between the reference site and study site provides a good indication of the effect of the tide restriction on tidal hydrology.



*Little River, North Hampton, NH: pre-restoration (above) and post-restoration (below).
Photos courtesy of NRCS*

There are two types of restrictive tidal crossings. One occurs when the opening of the culvert or bridge is too small or has started collapsing and does not allow natural amounts of water to pass through during each tidal cycle. The most common effect of this type of restriction is a decrease in salinity and especially flooding at high tide in the restricted marsh. The second type of restrictive crossing occurs where a culvert is elevated too high in relation to the creek bed. In this case, sufficient amounts of water may enter the restricted marsh during an incoming tide, but with a delayed effect since the tidal level in the unrestricted side must reach the height of the culvert before passing through it. Elevated culverts may prevent complete drainage of the restricted side because water cannot leave once water levels drop below the culvert, and even during low tide, there is standing water in the restricted marsh. Bank erosion may be evident on either side of the culvert with both types of tidal restrictions. Bank erosion resulting from tide restrictions is often described as “round-shaped pools,” which form on either side and directly next to the culvert.

Why Should Tidal Flow Be Monitored?

A reduction in tidal flow can have numerous adverse effects on salt marshes, the most important of which is a change in natural salinity regimes. Many plants and animals that exist in salt marshes are adapted to a specific range of physical and chemical conditions, and large-scale alterations such as tide restrictions can cause intolerant species to perish. When salinity levels fall below 20 ppt, the invasion of opportunistic brackish plants such as *Phragmites australis* becomes a problem. Tide restrictions may also block the passage of estuarine invertebrates and fish into the upper estuary, thereby reducing the export of organic matter from the salt marshes. A reduction in tidal flushing may result in the accumulation of detritus, nutrients, and pollutants in the restricted marsh.

How Should Tidal Flow Be Monitored?

Automatic water level recorders will be installed and operated by project coordinators for a minimum of two weeks, i.e., one lunar cycle of spring and neap tides (one month, or two lunar cycles is better) near the

source of tidal influx. For tidally restricted marshes, recorders will be installed both upstream and downstream of the tidal restriction. When automatic data collection gauges are unavailable, volunteers can be used to collect 10-minute measurements over 13-hour periods using a simple tide staff (a vertical ruler fixed in the tidal channel) for three spring and three neap tides would provide adequate information. Using either method, the elevations of the upstream and downstream devices are required, preferably referenced to National Geodetic Vertical Datum (NGVD).

Knowledge of the tidal current in the main channel can be useful when designating the tidal conduit for a tidal restoration and to assess the function of the current structure. Tidal current should be assessed over several tidal cycles and can be measured with a recording current meter.

Groundwater Levels

Healthy salt marshes depend on high groundwater levels to maintain pool and panne water depth. Groundwater drainage increases in proximity to manmade ditches, resulting in surface subsidence, soil compaction, and dry peat. These conditions can alter natural marsh hydrology, reduce plant productivity, and create opportunities for non-native invasive species to colonize the site.

Why Should Groundwater Levels Be Monitored?

Under circumstances where an important goal of the restoration project is to increase tidal flooding in order to reduce invasive plant species or increase open water habitat, water table depth monitoring is recommended.

Where Should Groundwater Levels Be Monitored?

Wells should be placed according to recommendations given for soil salinity stations along the upland edge of the marsh, or in sections of marsh potentially affected by the restoration action.

When Should Groundwater Levels Be Monitored?

Sampling will occur at low tide about six times a year in the early to mid growing season and include neap and spring tides.

How Should Groundwater Be Sampled?

Equipment

Piezometers are wells used to monitor groundwater levels. They are open only near the base to allow water into the well. They can be constructed from PVC pipe with a screened, perforated interval that intersects the water table. Changes in water table can be monitored with piezometers placed deep enough in the soil to intersect the water table during drier periods.

Sampling Methods

Various methods may be used for sampling groundwater. One of the methods you may use requires a dowel, the other a voltmeter. Both are valid techniques of measuring groundwater.

Dowel

The instrument used in this method is a dowel that measures one-meter in length and ___ centimeters in diameter and is graduated in centimeters. This size dowel is selected to displace the least amount of water.

1. Remove the well cap.
2. Using the dowel, measure the length of the well above the marsh surface by measuring from the ground to the top of the well (the outside of the well).
3. Slowly insert the dowel inside the well until the dowel hits the bottom of the well.
4. Remove the dowel and observe where the water level is on the dowel (the highest wet mark on the dowel). Record this number on the data sheet as inside height.

Voltmeter

1. Remove the well cap.
2. Turn the voltmeter dial to OHM.
3. SLOWLY lower the voltmeter inside the well, while observing the needle.
4. When the needle on the voltmeter starts to move, discontinue lowering the voltmeter and record the length from where you stopped lowering the voltmeter to the top of the well (this number will be in cm and should be recorded on the data sheet as inside height).

Appendix A. Suggested Reading

- Attenborough, D. 1998. *The Life of Birds*. Princeton University Press.
- Bertness, M.D. and A.M. Ellison. 1987. Determinants of Pattern in a New England Salt Marsh Plant Community. *Ecological Monographs* 57 (2): 129-147.
- Bertness, M.D. 1999. *The Ecology of Atlantic Shorelines*. Sinauer Associates, Inc. Sunderland, MA.
- Bieglow, H.B. and W.C. Schroder. 1953. *Fishes of the Gulf of Maine*. U.S. Fish and Wildlife Service, Fishery Bulletin 53 (74).
- Brinson, M.M. and R. Rheinhardt. 1996. The Role of Reference Wetlands in Functional Assessment and Mitigation. *Ecological Applications* 6 (1): 69-76.
- Brown, M.T., K. Brandt, and P. Adamus. 1990. Indicator Fact Sheets for Wetlands. In: *Ecological Indicators for the Environmental Monitoring and Assessment Program* (Hunsaker and Carpenter, eds.). US EPA, Office of Research and Development. EPA 600/3-90/060.
- Burdick, D., R. Buchsbaum, C. Cornelisen, and T. Diers. 1999. *Monitoring Restored and Created Salt Marshes in the Gulf of Maine: Framework and Data Collection Methods to Guide Monitoring Workshop*, June 2, 1998, Castle Hill, Ipswich, Massachusetts. Sponsored by: Massachusetts Audubon Society and Gulf of Maine Council on the Marine Environment.
- Carlisle, B.K., A.L. Hicks, J.P. Smith, S.R. Garcia, and B.G. Largay. 1998. *Wetland Ecological Integrity: An Assessment Approach: The Coastal Wetlands Ecosystem Protection Project*. MCZM, Boston, MA.
- Carlisle, B.K., A.L. Hicks, J.P. Smith, S.R. Garcia, and B.G. Largay. 1999. Plants and Aquatic Vertebrates as Indicators of Wetland Biological Integrity in Waquoit Bay Watershed, Cape Cod. *Environment Cape Cod* 2 (2): 30-60.
- Dates, G. and A. Reed. 2000. *Study Design: Deciding Why, What, How, When, and Where to Monitor*. Proceedings of the 6th National Volunteer Monitoring Conference, April 26-29, 2000. Austin, TX.
- Dates, G., A. Lehrer, J. Schoen, and R. McVoy. 1997. *Merrimack River Watershed Study Design Workbook*. Second Addition. The Volunteer Environmental Monitoring Network, Merrimack River Watershed Council, Lawrence, MA.
- Day, J.W., C.A.S. Hall, W.M. Kemp, and A. Yanez-Arancibia. 1989. *Estuarine Ecology*. John Wiley & Sons, Inc.
- Dionne, M., F.T. Short, and D.M. Burdick. 1999. Fish Utilization of Restored, Created, and Reference Salt-Marsh Habitat in the Gulf of Maine. Pages 384-404 in L. Benaka, editor. *Fish Habitat: Essential Fish Habitat and Rehabilitation*. American Fisheries Society, Symposium 22, Bethesda, MD.
- Ely, E. 1998. Monitoring Wetlands: Deciding What to Measure. *The Volunteer Monitor: The National Newsletter of Volunteer Water Quality Monitoring*. Vol 10, No. 1.
- Ely, E. 2000. Monitoring Massachusetts Marshes. *The Volunteer Monitor: The National Newsletter of Volunteer Water Quality Monitoring*. Vol 12, No. 2.
- Gleason, H.A. and A. Cronquist. 1991. *Manual of Vascular Plants of Northeastern United States and Adjacent Canada*. The New York Botanical Garden, Bronx, NY.
- Green, R.H. 1979. *Sampling Design and Statistical Methods for Environmental Biologists*. John Wiley, New York.
- Hayek, L.C. and M.A. Buzas. 1997. *Surveying Natural Populations*. Columbia University Press, New York.

- Hemmond, H.F. and J. Benoit. 1988. Cumulative Impacts on Water Quality Functions of Wetlands. *Environmental Management* 12 (5): 6639-6653.
- Hunt, M., A. Mayo, M. Brossman, and A. Markowitz. 1996. *The Volunteer Monitor's Guide to Quality Assurance Project Plans*. EPA 841-B-96-003. U.S. Environmental Protection Agency, Office of Wetlands, Oceans, and Watersheds. Washington, DC.
- Karr, J.R. and E.W. Chu. 1999. *Restoring Life in Running Waters: Better Biological Monitoring*. Island Press, Washington, DC.
- Kaufman, K. 1996. *Lives of North American Birds*. Houghton Mifflin Company.
- Kaufman, K. 2000. *Birds of North America*. Houghton Mifflin Company.
- Krebs, C.J. 1985. *Ecology: The Experimental Analysis of Distribution and Abundance*, 3rd Edition. Harper and Row, New York.
- Leibowitz, N.C. and M.T. Brown. 1990. Indicator Strategy for Wetlands. In: *Ecological Indicators for the Environmental Monitoring and Assessment Program*. Hunsaker and Carpenter (Eds.) US EPA, Office of Research and Development. EPA 600/3-90/060.
- Mitch, W.J. and J.G. Gosselink. 1993. *Wetlands*. Van Nostrand Reinhold, Inc. New York, NY.
- Murdy, E.O., R.S. Birdsong, and J.A. Musick. 1997. *Fishes of Chesapeake Bay*. Smithsonian Institute.
- National Geographic Society. 1999. *Field Guide to the Birds of North America*, 3rd Edition. National Geographic Society.
- National Geographic Society. 1986. *Guide to Bird Songs*. National Geographic Society and the Cornell University Library of Natural Sounds.
- Natural Resources Conservation Service and Partners. 2001. *Evaluation of Restorable Salt Marshes in New Hampshire*. http://www.nh.nrcs.usda.gov/Ecosystem_Restoration/
- Neckles, H.A. and M. Dionne. 2000. *Regional Standards to Identify and Evaluate Tidal Wetland Restoration in the Gulf of Maine*. Wells National Estuarine Research Reserve Technical Report, Wells ME. http://restoration.nos.noaa.gov/htmls/resources/general_pubs.html#N
- New Hampshire Department of Environmental Services, Coastal Program. 2003. *Life in New Hampshire Salt Marshes: A Quick-Reference Field Guide*. Portsmouth, NH.
- Niedowski, N.L. 2000. *New York State Salt Marsh Restoration and Monitoring Guidelines*. New York State Department of State, Division of Coastal Resources, Albany, NY and New York State Department of Environmental Conservation, East Setauket, NY.
- Niering, W.A. and R.S. Warren. 1980. Vegetation Patterns and Processes in New England Salt Marshes. *Bioscience* 30 (5): 301-307.
- Ott, R.L. 1993. *An Introduction to Statistical Methods and Data Analysis*. Duxbury Press, California.
- Peterson, R.T. 1980. *Field Guide to Birds East of the Rockies*. Houghton Mifflin Company.
- Peterson, R.T. 1999. *Field Guide to Bird Songs – Eastern/Central North America*. Cornell University Library of Natural Sounds.

- Pollock, L.W. 1998. *A Practical Guide to the Marine Animals of Northeastern North America*. Rutgers University Press.
- Purinton, T.A. and D.C. Mountain. 1998. *Tidal Crossing Handbook: A Volunteer Guide to Assessing Tidal Restrictions*. Parker River Clean Water Association, Byfield, MA.
- Robins, C.R. and G.C. Ray. 1986. *A Field Guide to Atlantic Coast Fishes (North America)*. The Peterson Field Guide Series. Houghton Mifflin Company, Boston, MA.
- Rozas, L.P. and T.J. Minello. 1997. Estimating Densities of Small Fishes and Decapod Crustaceans in Shallow Estuarine Habitats: A Review of Sampling Design With Focus on Gear Selection. *Estuaries* 20 (1): 199-213.
- Rozas, L.P. 1995. Hydroperiod and Its Influence on Nekton Use of the Salt Marsh: A Pulsing Ecosystem. *Estuaries* 18 (4): 579-590.
- Saltonstall, K. 2002. Cryptic Invasion By a Non-Native Genotype of the Common Reed, *Phragmites australis*, Into North America. *Proceedings of the National Academy of Sciences* 99 (4): 2445-2449.
- Schreck, C.B. and P.B. Moyle. 1990. *Methods for Fish Biology*. American Fisheries Society, Bethesda, MD.
- Sibley, D.A. 2000. *National Audubon Society Sibley Guide to Birds*. Alfred A. Knopf, Inc.
- Sibley, D.A. 2001. *The Sibley Guide to Bird Life and Behavior*. Alfred A. Knopf, Inc.
- Sinicrope, T.L., G. Hine, R.S. Warren, and W.A. Niering. 1990. Restoration of an Impounded Salt Marsh in New England. *Estuaries* 13 (1): 25-30.
- Stokes, S. and L. Stokes. 1996. *Stokes Field Guide to Birds: Eastern Region*. Little Brown and Company.
- Teal, J. and M. Teal. 1969. *Life and Death of the Salt Marsh*. Ballentine Books, New York, NY.
- Tiner, R.W. 1987. *A Field Guide to Coastal Wetland Plants of the Northeastern United States*. The University of Massachusetts Press, Amherst MA.
- Tiner, R.W. 1993. *Coastal Wetland Plants of the Southeastern United States*. The University of Massachusetts Press, Amherst MA.
- United States Environmental Protection Agency. 1993. *Volunteer Estuary Monitoring: A Methods Manual*. EPA 842-B-93-004. U.S. Environmental Protection Agency, Office of Water, Washington, D.C.
- Walton, R.K. and R.W. Lawson. 1994. *Peterson Field Guides: Eastern and Central More Birding by Ear*. Cornell University Library of Natural Sounds.
- Walton, R.K. and R.W. Lawson. 1999. *Peterson Field Guides: Eastern/Central Birding by Ear*. Cornell University Library of Natural Sounds.
- Waters, W.E. and D.C. Erman. 1990. Research Methods: Concept and Design. Pages 1-34 in: C.B. Shreck and P.B. Moyles, eds. *Methods for Fish Biology*. American Fisheries Society, Bethesda, MD.
- Weiss. H.M. 1995. Marine Animals of Southern New England and New York: Identification Keys to Common Nearshore and Shallow Water Macrofauna. State Geological and Natural History Survey of Connecticut, Bulletin 115.

Appendix B. Glossary

Abundance: The amount — by count, weight, or other measure — of a given group in a given area. Generally, abundance refers to the number of individuals of a species (genus, family) within an area of survey.

Alga/Algae: Very simple, often one celled, plants that are either attached or unattached in aquatic (marine or freshwater) environments; can be used as a term to cover simple seaweed. Floating clumps of algae are called alga mats.

Ambient Water Quality: The conditions of a water body (or wetland) generally taken as a whole (e.g. the average pH of Pleasant Bay in 1999), contrasted with site source specific or episodic measurements.

Biological Impairment: Diminished quality, strength, or value of the condition of an individual, group, habitat, and/or function of living organisms.

Biological Integrity: Ability of an ecosystem to support and maintain a balanced, adaptive community of organisms having a species composition, diversity, and functional organization comparable to that of pristine habitats within a region.

Community: A group of species inhabiting a given area, where organisms interact and influence one another's distribution, abundance, and evolution.

Detritus: Dead and decomposing plant and animal material.

Diversity: Variety or heterogeneity in taxonomic groups.

Environmental Stressor: Any material or process (physical, chemical, or biological) that can adversely affect a salt marsh, includes both natural and human disturbances.

Estuary: Region of interaction between rivers and near-shore ocean waters, where tidal action and river flow mix fresh and salt water. Such areas include bays, mouths of rivers, salt marshes, and lagoons. These brackish water ecosystems shelter and feed marine life, birds, and wildlife.

Eutrophication: The process by which a body of water becomes enriched with nutrients, particularly nitrogen and phosphorus, which usually changes ecosystem properties and functions.

Evaluation: Area A comparable and representative portion of a reference or study salt marsh.

Food Web: The linkage of organisms based on their feeding relationships sources.

Generalist: A species (genus, family) that is able to exist or thrive in a variety of habitats or conditions.

Global Positioning System (GPS): Technology that utilizes communication between orbiting satellites and ground receivers to pinpoint exact locations on the earth.

Groundwater: The water found beneath the Earth surface, frequently used in reference to aquifers and drinking water wells.

Habitat: The sum of the physical, chemical, and biological environment occupied by individuals or a particular species, population, or community.

High Marsh: The area of a New England salt marsh that is flooded by higher than average tides and dominated by the grasses *Spartina patens* and *Distichlis spicata*. The high marsh lies between the low marsh and the marsh's upland border.

Human Disturbance: Activity or state caused, directly or indirectly, by humans that intrudes, interrupts, or

perturbs the natural state of ecological relationship and function.

Hydrology: The [study of] water of the earth, its occurrences, distribution, and circulation with particular emphasis on the chemistry and movement of water.

Indicator: An attribute or measure that is strongly suggestive of the condition or direction of an ecological system.

Introduced Species: See non-indigenous.

Invasive Species: Non-indigenous organisms that may threaten the diversity or abundance of native species or natural ecological relationships and functions by spreading and out-competing native species.

Invertebrate(s): Animals without internal skeletons and backbones. Marine invertebrates live in ocean-derived salt water, freshwater invertebrates live in freshwater for at least part of their life cycle, and terrestrial invertebrates are associated with uplands and fringes of aquatic habitats.

Low Marsh: The seaward area of a salt marsh, generally flooded daily by the tides, and dominated by the tall form of *Spartina alterniflora*.

Macroinvertebrate: An animal without an internal backbone that is large enough to be seen by the naked eye.

Marsh Border: The zone of a salt marsh that is only flooded during extreme high tides or coastal storms, and sustains a variety of upland and wetland plants that are not well adapted to periodic flooding or salt stress.

Monitoring: Periodic or continuous survey or sampling to determine the status or condition of various media and systems, including water bodies, groups of plants and animals, or ecological systems.

Morphology: The [study of] form and structure of an organism.

Nekton: Any organisms that actively swim in the water column.

Non-Indigenous: A species transported intentionally or accidentally from another region, allowing it to occur in areas beyond its normal range. Synonym: introduced species.

Panne: A depression on the surface of a salt marsh. This term is used variably in the literature and field to include both vegetated and un-vegetated, as well as permanently or temporarily flooded depressions.

Parameter: A measurable property whose value determines characteristics of an ecosystem (e.g. salinity is a measurable attribute of estuarine waters).

Plot Sample: A field sample technique that gathers information from an area enclosed within the dimensions set by a frame of a standard size.

Pool: A depression on the surface of a salt marsh generally permanently flooded depressions.

Population: A group of interbreeding organisms occupying a particular space or area; all of the organisms that constitute a specific group or occur in a specified habitat.

Pore Water: The shallow groundwater occupying the interstitial areas (or pores) of marsh substrate.

Quadrat Sample: See plot sample.

Qualitative: Involving distinctions based on standards, traits, or value.

Quantitative: Expressible as, or relating to, a measurable value.

Reference Marsh/Site: A marsh that exhibits a typical “minimally disturbed” condition, or maximum functional

capacity; and represents other marshes in a specific region sharing the same water regime, topographic setting, and climate zone.

Refractometer: A device used to measure salinity (or the concentrations of certain dissolved minerals). Prisms send light through a very small water sample and the bend of the light is consistent with the concentration (amount) of salts.

Rhizomes: A horizontal, usually underground stem that generally sprouts roots and shoots from its nodes.

Salinity Regime: The measured, normal fluctuations in salinity over tidal and seasonal cycles.

Salt Marsh: Low-lying, vegetated coastal wetlands, influenced by the tidal estuary or marine waters.

Sample Station: A specific location within the salt marsh site selected to conduct field sampling.

Sensitive: Organisms that have a low tolerance of pollution and disturbance. Numbers tend to decrease with impact.

Spawning: The release of gametes or eggs into the water.

Specialist: An organism with very specific requirements for some aspects of its ecology or phases of its life cycle.

Substratum: The various materials that collectively make up the exposed or submerged surfaces of wetlands and aquatic environments, which may include sand, silt, peat, algae, logs, wood, debris, bank surface, sediments, leaf packs, mud, rock, and sometimes solid waste such as tires.

Taxonomy: The study of the relationships and classification of organisms.

Tidal Restriction: A structure or landform that restricts natural tidal flow, such as a culvert, bridge, dam, or causeway.

Tolerant: Organisms that have a high tolerance of pollution or disturbance, whose numbers tend to increase with impact.

Transect: A method for environmental sample or survey using a straight line to delineate the area of analysis.

Vagile: A species that easily disperses over large distances.


Wetland: Areas where water covers the soil, or is present either at or near the surface of the soil for at least part of the growing season.

Zonation: The observed occurrence of New England salt marsh plants to organize into apparently discrete areas, due to flooding, salinity, and other forcing factors. A classic zonation pattern is (progressing across a marsh from estuarine water to the upland) low marsh, high marsh, and border or fringing marsh. In many marshes, the classic pattern does not hold, and the plant communities would be better described as a patchwork or mosaic (see Figure 5.1).

Appendix C. Lift Net & Ditch Net Materials and Construction

Materials for Ditch Net Construction (For 1 Net)

- NYLON netting (24 lb test), 1/8" mesh, at least in deep. Each net takes 5 m of netting - a 1 m x 3 m section for the center of the net (sides & bottom) and two 1 m sections for the doors
- 20 m of nylon rope, 3/16" diameter. Each net takes 20 m of line = four 4 m lengths for rip cords and four in lengths for runner lines of the doors
- 5 m of leadcore line; m for the top of each door (total 2 m) and 3 m for the floor of the net
- 4 eye-hooks
- 4 oak stakes — 1.5 to 2 m long, 2.5 cm square
- Staple gun and 3/8" stainless steel staples
- D-ring hand pliers and 9/16" C-ring fasteners
- 25 to 30 plastic rings or rubber O-rings, approximately 2.5 cm diameter



Ditch Net Construction

- Cut a 1 m by 3 m section of the nylon netting for the center of the net.
- Cut two 1 m by 1 m sections of nylon netting for the doors of the net.
- Attach the doors of the net to the center section. The doors should be centered on the main body of the net along the 3 m length (Fig 1). To attach the doors take a 1 m length of leadcore line and wrap the nylon netting from the leading edge of the door and the center 1 m middle section of the net body around the leadcore and fasten the two pieces of nylon netting to the leadcore line with the D-ring pliers and 9/16" C-ring fasteners.
- Attach 5 to 7 nylon rings or rubber O-rings to sides of the doors (side A in Fig 1). Use the D-ring pliers to attach the rings to the nylon netting. The rings should be attached to the edge of the netting so the center of the ring is clear of the netting. The draw cord that pulls the doors up passes through these rings.

Attach 3 to 5 plastic rings to the top of the door (side B in Fig. 1). Use the D-ring pliers to attach the rings to the nylon netting. The rings should be attached to the edge of the netting so the center of the ring is clear of the netting.

- Attach a short length of lead core line to the top of each door (Fig. 1, side B) using either cable ties or the D-ring pliers and C-ring fasteners. This is to weigh down the top of the net so it does not float up, and impede the passage of fish through the net.

Attach a length (approximately 1 m) of leadcore line to the bottom center of the net (Fig. 1b) on the outside of the net using either cable ties or the D-ring pliers and C-ring fasteners. This is to weigh down the center of the net so it does not float up when placed in the ditch.

- Attach the net to the four oak stakes using a staple gun and stainless steel staples. The free edges of the net (Fig 1, side C, and Fig. 1 between points 1 and 2) are stapled to the oak stakes. The portion of the net closest to the doors should be stapled starting at approximately 30 cm (1 ft) from the bottom of the oak stake, and continue up towards the top of the stake. The bottom 1 ft of the stake should be free of the net so that the stake can be pushed into the ground to hold the net in place while it is deployed.
- The runner lines are attached next. The runner lines hold the plastic rings close to the stake, so when the door is pulled up the net remains close to the stake.
- Attach the bottom of runner lines to the interior of the stakes (on top of the stapled netting). The runner lines are approximately 1 m in length. The bottom of the runner line should be attached at the intersection of the doors and main body of the net. Tie a few knots in the end of the line and staple the line to the stakes using several staples close together on each side of the knot so the line will not pull loose.

Pass the free end of the runner line through the 5 plastic rings that are attached to side A (Fig. 1) of the door closest to the stake (Fig. 1, runner line (3) and plastic rings (4)). The bottom most ring is added first, then the next ring, until all rings for that door side are on the runner line. The runner line is then pulled taut against the stake and the free end is stapled approximately 5 to 8 cm above the end of the net. After stapling a knot should be tied in the free end of the line and stapled again on either side of the knot to ensure the runner line does not come loose.

- Attach the ripcord to the center ring on the top of the door, and pass the ripcord through one of the rings on the corner of the door. Then pass the ripcord through the top ring of the door that is attached to the runner line. Attach another ripcord to the same center ring, and pass it through the other corner ring, and the top ring on the other side of the door. When these lines are pulled, they will pull on the top rings attached to the doors, which in turn will pull the sides of the doors up the stakes to enclose the sides of the net.
- Attach the ripcords to the other side of the net as described above.


Attach the eyehook to the oak stake. When the net is held upright, with the 4 stakes sticking into the ground, the eyehook should be placed on the outside of the stake. The free end of ripcord is passed through the eyehook. When the ripcord is pulled the line should pass easily through the eyehook, so the doors are pulled up smoothly.

Label the stakes. We usually label the stakes A, B, C, and D. Be sure to label each net exactly the same. The labels are used to set the net correctly in the ditch and to measure the distance between the stakes in

Materials for Lift Net Construction

- Nylon netting (24 lb test), 1/8" mesh.
- 3/16" diameter nylon rope
- Size 50 lead-line
- 9/16" Ringer
- 9/16" C ring fasteners
- Large sewing needle
- Spool of sturdy thread – purchased at hardware store.

Lift Net Construction



Lift Net Construction

- Cut a piece of nylon netting measuring 4 ft wide and 8.5 ft long (some will be used in the sewing so your total length will be a little less when completed).
- Cut two 28-30 ft pieces 3/16" diameter nylon rope.
- Cut one 17 ft piece of size 50 leadline.
- To ensure it will not unravel with time, burn the ends of the leadline and while it is hot, flatten the end with a wrench.
- Find the middle of the nylon rope and mark it with a pen
- Line up the rope along one end of the nylon net -along the width side (see Fig 1.)
- Bring rope up from edge of netting 1/2-1 inch placing the mark that you made on the rope to the middle of the net.
- Starting from middle of net (this makes it easier) start sewing the rope onto the netting. After reaching the end, double back and redo the same edge to ensure that the rope is securely attached to the netting. Complete the other half of the edge. Repeat for other side.
- Cut off extra netting if necessary.
- After both edges of netting are attached to rope, the leadline needs to be attached to the bottom of the net using a ringer. Usually, this is simply done by making a large S in the middle of your net using your leadline and attaching it with a ringer.
- It may be easier to lay your leadline down first and then place the nylon net over it, adjust the leadline to the position you would like it to be and then use the ringer to attach it. This is a much easier way of using the ringer. The locations in which to ring the rope are shown in Fig 2.

Appendix D: Data Sheets